

US009526764B2

(12) **United States Patent**
Werner et al.

(10) **Patent No.: US 9,526,764 B2**
(45) **Date of Patent: Dec. 27, 2016**

(54) **COMBINATION OF AN INSULIN AND A GLP-1-AGONIST**

(75) Inventors: **Ulrich Werner**, Frankfurt am Main (DE); **Barbel Rotthäuser**, Frankfurt am Main (DE)

(73) Assignee: **SANOFI-AVENTIS DEUTSCHLAND GMBH**, Frankfurt am Main (DE)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/123,835**

(22) PCT Filed: **Oct. 9, 2009**

(86) PCT No.: **PCT/EP2009/063195**
§ 371 (c)(1),
(2), (4) Date: **Sep. 30, 2011**

(87) PCT Pub. No.: **WO2010/043566**
PCT Pub. Date: **Apr. 22, 2010**

(65) **Prior Publication Data**
US 2012/0021978 A1 Jan. 26, 2012

(30) **Foreign Application Priority Data**

Oct. 17, 2008 (DE) 10 2008 051 834
Oct. 24, 2008 (DE) 10 2008 053 048
Aug. 20, 2009 (DE) 10 2009 038 210

(51) **Int. Cl.**
A61K 38/28 (2006.01)
A61K 38/00 (2006.01)
A61K 38/26 (2006.01)

(52) **U.S. Cl.**
CPC **A61K 38/28** (2013.01); **A61K 38/26** (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,758,683 A 9/1973 Jackson
3,868,358 A 2/1975 Jackson
4,153,689 A 5/1979 Hirai et al.
4,258,134 A 3/1981 Yoshida et al.
4,367,737 A * 1/1983 Kozam et al. 604/191
4,608,364 A 8/1986 Grau
4,614,730 A 9/1986 Hansen et al.
4,644,057 A 2/1987 Bicker et al.
4,689,042 A 8/1987 Sarnoff et al.
4,701,440 A 10/1987 Grau
4,731,405 A 3/1988 Kirsch et al.
4,783,441 A 11/1988 Thurow

4,839,341 A 6/1989 Massey et al.
4,863,902 A 9/1989 Amagase et al.
4,885,164 A 12/1989 Thurow
4,923,162 A 5/1990 Fleming et al.
4,959,351 A 9/1990 Grau
4,960,702 A 10/1990 Rice et al.
4,994,439 A 2/1991 Longenecker et al.
5,008,241 A 4/1991 Markussen et al.
5,034,415 A 7/1991 Rubin
5,070,186 A 12/1991 Joergensen
5,101,013 A 3/1992 Dorschug et al.
5,177,058 A 1/1993 Dorschug
5,187,177 A 2/1993 Garzaran
5,227,293 A 7/1993 Stengelin et al.
5,253,785 A 10/1993 Haber et al.
5,272,135 A 12/1993 Takruri
5,358,708 A 10/1994 Patel
5,358,857 A 10/1994 Stengelin et al.
5,370,629 A 12/1994 Michel et al.
5,397,771 A 3/1995 Bechgaard et al.
5,407,609 A 4/1995 Tice et al.
5,424,286 A 6/1995 Eng
5,428,006 A 6/1995 Bechgaard et al.
5,473,049 A 12/1995 Obermeier et al.
5,474,978 A 12/1995 Bakaysa et al.
5,478,323 A 12/1995 Westwood et al.
5,496,924 A 3/1996 Habermann et al.
5,506,203 A 4/1996 Backstrom et al.
5,509,905 A 4/1996 Michel
5,514,646 A 5/1996 Chance et al.
5,524,286 A 6/1996 Chiesa et al.
5,534,488 A 7/1996 Hoffmann
5,545,618 A 8/1996 Buckley et al.
5,547,929 A 8/1996 Anderson, Jr. et al.
5,559,094 A 9/1996 Brems et al.
5,595,756 A 1/1997 Bally et al.
5,597,796 A 1/1997 Brange
5,614,219 A 3/1997 Wunderlich et al.
5,614,492 A 3/1997 Habener

(Continued)

FOREIGN PATENT DOCUMENTS

AU 593274 B2 2/1990
AU 612324 B2 7/1991

(Continued)

OTHER PUBLICATIONS

Tews et al. 2007. Diabetes. 56(suppl. 1):A-72-A73.* Tews et al. 2008. Hormone and Metabolic Res. 40:172-80.*
www.fda.gov/CombinationProducts/AboutCombinationProducts/ucm101496.1, 2009, downloaded Jul. 13, 2012.*
Eng et al. 1992. J. Biol. Chem. 267:7402-7405.*
2012. WHO Rational Use of Medicines, http://www.who.int/medicines/areas/rational_use/en/ downloaded Dec. 18, 2014 10:02:48 AM.*
<http://web.archive.org/web/20081021043832/http://www.rxlist.com/lantus-drug.htm>, downloaded Nov. 12, 2015.*
<http://www.pipelinereview.com/>, published Jun. 9, 2008, downloaded Nov. 12, 2015.*

(Continued)

Primary Examiner — Shulamith H Shafer
(74) *Attorney, Agent, or Firm* — McDonnell Boehnen Hulbert & Berghoff LLP

(57) **ABSTRACT**

The invention relates to a drug comprising at least one insulin and at least one GLP-1 receptor agonist.

15 Claims, 10 Drawing Sheets

US 9,526,764 B2

Page 2

(56)

References Cited**U.S. PATENT DOCUMENTS**

5,631,224 A	5/1997	Efendic et al.	8,084,420 B2	12/2011	Steiner et al.
5,654,008 A	8/1997	Herbert et al.	8,092,421 B2	1/2012	Seiferlein et al.
5,656,722 A	8/1997	Dorschug	8,092,422 B2	1/2012	Seiferlein et al.
5,663,291 A	9/1997	Obermeier et al.	8,178,495 B2 *	5/2012	Chilkoti 514/11.7
5,670,360 A	9/1997	Thorens	8,574,214 B2	11/2013	Kuehn et al.
5,693,608 A	12/1997	Bechgaard et al.	8,633,156 B2	1/2014	Habermann et al.
5,700,662 A	12/1997	Chance et al.	8,735,349 B2	5/2014	Silvestre et al.
5,707,641 A	1/1998	Gertner et al.	2001/0012829 A1	8/2001	Anderson et al.
5,783,556 A	7/1998	Clark et al.	2001/0033868 A1	10/2001	Rossling et al.
5,824,638 A	10/1998	Burnside et al.	2001/0039260 A1	11/2001	Havelund
5,846,747 A	12/1998	Thorens et al.	2001/0047084 A1	11/2001	Knudsen et al.
5,846,937 A	12/1998	Drucker	2002/0107265 A1	8/2002	Chen et al.
5,948,751 A	9/1999	Kimer et al.	2002/0132760 A1	9/2002	Van Antwerp et al.
5,952,297 A	9/1999	De Felippis et al.	2002/0177151 A1	11/2002	Gimeno
5,985,309 A	11/1999	Edwards et al.	2002/0198140 A1	12/2002	Havelund
6,006,753 A	12/1999	Efendic	2003/0004096 A1	1/2003	Boderke
6,034,054 A	3/2000	DeFelippis et al.	2003/0026872 A1	2/2003	Dake et al.
6,043,214 A	3/2000	Jensen et al.	2003/0104983 A1	6/2003	DeFelippis et al.
6,051,551 A	4/2000	Hughes et al.	2003/0170691 A1	9/2003	Gimeno et al.
6,051,689 A	4/2000	Thorens	2004/0048783 A1	3/2004	Brunner-Schwarz et al.
6,100,376 A	8/2000	Dorschug	2004/0092590 A1	5/2004	Arterburn et al.
6,110,703 A	8/2000	Egel-Mitani et al.	2004/0097410 A1	5/2004	Zheng et al.
6,174,856 B1	1/2001	Langballe et al.	2004/0106547 A1	6/2004	Larsen et al.
6,191,102 B1	2/2001	DiMarchi et al.	2004/0186046 A1	9/2004	Burgess et al.
6,211,144 B1	4/2001	Havelund	2004/0229774 A1	11/2004	Rosskamp et al.
6,227,819 B1	5/2001	Gettel et al.	2004/0235710 A1	11/2004	DeFelippis et al.
6,248,363 B1	6/2001	Patel et al.	2005/0014679 A1	1/2005	Beals et al.
6,267,981 B1	7/2001	Okamoto et al.	2005/0079996 A1	4/2005	Horiguchi et al.
6,268,335 B1	7/2001	Brader	2005/0106147 A1	5/2005	Jordan et al.
6,268,343 B1	7/2001	Knudsen et al.	2005/0171009 A1	8/2005	Brunner-Schwarz et al.
6,271,241 B1	8/2001	DeSimone et al.	2005/0209142 A1	9/2005	Bertilsson et al.
6,284,725 B1	9/2001	Coolidge et al.	2006/0004049 A1 *	1/2006	Yao et al. 514/317
6,309,663 B1	10/2001	Patel et al.	2006/0014678 A1	1/2006	Cowley et al.
6,310,038 B1	10/2001	Havelund	2006/0019347 A1	1/2006	Cho et al.
6,329,336 B1	12/2001	Bridon et al.	2006/0057137 A1	3/2006	Steiness
6,335,316 B1	1/2002	Hughes et al.	2006/0073213 A1	4/2006	Hotamisligil et al.
6,344,180 B1	2/2002	Holst et al.	2006/0093576 A1	5/2006	Chen et al.
6,358,924 B1	3/2002	Hoffmann	2006/0194719 A1	8/2006	Ebbelohj et al.
6,384,016 B1	5/2002	Kaarsholm	2006/0239933 A1	10/2006	Nilsson et al.
6,388,053 B1	5/2002	Galloway et al.	2006/0287221 A1	12/2006	Knudsen et al.
6,395,767 B2	5/2002	Robl et al.	2007/0027063 A1	2/2007	Boss et al.
6,410,508 B1	6/2002	Isales et al.	2007/0111940 A1	5/2007	Larsen et al.
6,417,164 B1	7/2002	Kolterman et al.	2007/0128193 A1	6/2007	O'Neil et al.
6,444,641 B1	9/2002	Flora	2007/0135338 A1	6/2007	O'Neil et al.
6,468,959 B1	10/2002	Wunderlich et al.	2007/0155653 A1	7/2007	Boderke
6,489,292 B1	12/2002	Havelund et al.	2007/0191271 A1	8/2007	Mayhew et al.
6,528,486 B1	3/2003	Larsen et al.	2007/0237827 A1	10/2007	Sung et al.
6,734,162 B2	5/2004	Van Antwerp et al.	2008/0064856 A1	3/2008	Warne et al.
6,767,887 B1	7/2004	Hoffmann et al.	2008/0146490 A1	6/2008	Joabsson et al.
6,818,738 B2	11/2004	Havelund	2008/0234200 A1	9/2008	Quay et al.
6,852,694 B2	2/2005	Van Antwerp et al.	2008/0248999 A1	10/2008	Steiner
6,875,589 B1	4/2005	Doerschug et al.	2008/0260840 A1	10/2008	Alessi et al.
6,908,610 B1	6/2005	Sato	2008/0267907 A1	10/2008	Poulsen
6,908,897 B2	6/2005	Brandenburg et al.	2009/0082255 A1	3/2009	Brunner-Schwarz et al.
6,960,561 B2	11/2005	Boderke	2009/0088369 A1	4/2009	Steiness
6,969,702 B2	11/2005	Bertilsson et al.	2009/0099064 A1	4/2009	Lougheed
7,022,674 B2	4/2006	DeFelippis et al.	2009/0175840 A1	7/2009	Kashyap et al.
7,115,563 B2	10/2006	Younis et al.	2009/0176692 A1	7/2009	Habermann et al.
7,119,086 B2	10/2006	Di Malta et al.	2009/0180953 A1	7/2009	Gotthardt et al.
7,192,919 B2	3/2007	Tzannis et al.	2009/0186819 A1	7/2009	Carrier et al.
7,205,276 B2	4/2007	Boderke	2009/0208565 A1	8/2009	Ebbelohj et al.
7,205,277 B2	4/2007	Boderke	2009/0214657 A1	8/2009	Qazi et al.
7,238,663 B2	7/2007	DeFelippis et al.	2009/0304665 A1	12/2009	Frost et al.
7,405,196 B2	7/2008	Rosskamp et al.	2009/0312236 A1	12/2009	Beals et al.
7,476,652 B2	1/2009	Brunner-Schwarz et al.	2009/0324701 A1	12/2009	Williams
7,544,656 B2	6/2009	Sabetsky	2010/0029558 A1	2/2010	Bristow
7,544,657 B2	6/2009	Ebbelohj et al.	2010/0055049 A1	3/2010	Kuo et al.
7,576,050 B2	8/2009	Greig et al.	2010/0057194 A1	3/2010	Ryan
7,713,930 B2	5/2010	Brunner-Schwarz et al.	2010/0069292 A1	3/2010	Pohl et al.
7,803,763 B2	9/2010	Thurrow et al.	2010/0069293 A1	3/2010	Bolotin et al.
7,807,242 B2	10/2010	Soerensen et al.	2010/0227816 A1	9/2010	Flatt et al.
7,918,833 B2	4/2011	Veasey et al.	2010/0279931 A1	11/2010	Garibay et al.
7,939,293 B2	5/2011	Habermann et al.	2010/0311112 A1	12/2010	Rissom et al.
7,977,310 B2	7/2011	Rosskamp et al.	2011/0020294 A1	1/2011	Hammerman
8,048,854 B2	11/2011	Habermann et al.	2011/0021423 A1	1/2011	Olsen et al.
			2011/0077197 A1	3/2011	Habermann et al.
			2011/0118178 A1	5/2011	Silvestre et al.
			2011/0118180 A1	5/2011	Silvestre et al.
			2011/0144008 A1	6/2011	Larsen et al.

US 9,526,764 B2

(56)

References Cited

U.S. PATENT DOCUMENTS

2011/0152185	A1	6/2011	Plum et al.
2011/0173722	A1	7/2011	Habermann et al.
2011/0230402	A1	9/2011	Johansen et al.
2011/0236925	A1	9/2011	Hazra et al.
2011/0245165	A1	10/2011	Larsen et al.
2011/0281790	A1	11/2011	Pohl et al.
2011/0301081	A1	12/2011	Becker et al.
2012/0021978	A1	1/2012	Werner et al.
2012/0121611	A1	5/2012	Lodie et al.
2012/0122774	A1	5/2012	Becker et al.
2012/0183616	A1	7/2012	Sprogoe et al.
2012/0232002	A1	9/2012	Schoettle et al.
2012/0241356	A1	9/2012	Pfenninger et al.
2012/0252724	A1	10/2012	Schoettle et al.
2012/0277147	A1	11/2012	Boka et al.
2012/0283179	A1	11/2012	Brunner-Schwarz et al.
2012/0295846	A1	11/2012	Hagendorf et al.
2012/0316108	A1	12/2012	Chen et al.
2013/0005649	A1	1/2013	Niemoeller et al.
2013/0012433	A1	1/2013	Roskamp et al.
2013/0023467	A1	1/2013	Silvestre et al.
2013/0040878	A1	2/2013	Silvestre et al.
2013/0065828	A1	3/2013	Ruus et al.
2013/0079279	A1	3/2013	Becker et al.
2013/0085102	A1	4/2013	Silvestre et al.
2013/0096059	A1	4/2013	Stechl et al.
2013/0096060	A1	4/2013	Stechl et al.
2013/0116179	A1	5/2013	Hess et al.
2013/0203666	A1	8/2013	Niemoeller et al.
2013/0284912	A1	10/2013	Vogel et al.
2013/0296236	A1	11/2013	Silvestre et al.
2014/0148384	A1	5/2014	Boka et al.
2014/0206611	A1	7/2014	Becker et al.
2014/0221285	A1	8/2014	Bley et al.
2014/0248365	A1	9/2014	Rademacher et al.
2014/0371141	A1	12/2014	Souhami et al.

FOREIGN PATENT DOCUMENTS

CA	1173388	A	8/1984
CA	1258427	A	8/1989
CA	1336329	C	7/1995
CA	1341203	C	3/2001
CA	2662084	A1	3/2008
CN	1276731	A	12/2000
CN	1413582	A	4/2003
CN	101366692	A	2/2009
CN	101444618	A	6/2009
CN	101454019	A	6/2009
CN	101670096	A	3/2010
DE	2219635	A1	11/1972
DE	3240177	A1	5/1983
DE	19637230	A1	3/1998
DE	10 2008 003 566	A1	7/2009
DE	102008003568	A1	7/2009
DE	102008053048	A1	4/2010
EA	006019	B1	8/2005
EP	0018609	B1	9/1983
EP	0046979	B1	9/1983
EP	0132769	A1	2/1985
EP	0140084	A1	5/1985
EP	0166529	A1	1/1986
EP	0180920	A2	5/1986
EP	0200383	A2	11/1986
EP	0211299	A2	2/1987
EP	0214826	B1	3/1987
EP	0224885	A1	6/1987
EP	0227938	A2	7/1987
EP	0229956	A1	7/1987
EP	0229998	A2	7/1987
EP	0254516	A2	1/1988
EP	0305760	A2	3/1989
EP	0368187	A2	5/1990
EP	0375437	B1	6/1990
EP	0383472	A2	8/1990

EP	0419504	B1	4/1991
EP	0194864	B1	6/1992
EP	0600372	A1	6/1994
EP	0668282	A1	8/1995
EP	0668292	A2	8/1995
EP	0678522	B1	10/1995
EP	0837072	A2	4/1998
EP	0845265	A1	6/1998
EP	0885961	B1	12/1998
EP	1076066	A1	2/2001
EP	1172114	A2	1/2002
EP	1196444	A1	4/2002
EP	1222207	A1	7/2002
EP	1 523 993	A1	4/2005
EP	1364029	B1	12/2005
EP	1906991	A2	4/2008
EP	2112161	A2	10/2009
EP	2187950	A0	5/2010
EP	2324853	A1	5/2011
EP	2329848	A1	6/2011
EP	2389945	A1	11/2011
EP	0921812	B2	12/2011
EP	2387989	B1	7/2014
FR	2456522	A1	12/1980
GB	835638	A	5/1960
GB	840870	A	7/1960
GB	1527605	A	10/1978
GB	1554157	A	10/1979
JP	S61212598	A	9/1986
JP	S6399096	A	4/1988
JP	H02218696	A	8/1990
JP	2-264798	A	10/1990
JP	H03504240	A	9/1991
JP	H06506444	A	7/1994
JP	2001521004	A	11/2001
JP	2002516880	A	6/2002
JP	2003505347	A	2/2003
JP	2005508895	A	4/2005
JP	2005532365	A	10/2005
JP	2006-515267	A	5/2006
JP	2006137678	A	6/2006
JP	2007204498	A	8/2007
JP	2009091363	A	4/2009
JP	2009519961	A	5/2009
JP	2012505852	A	3/2012
JP	2012255040	A	12/2012
RU	2008116057	A	10/2009
RU	2386631	C2	4/2010
TW	157005	B	5/1991
TW	562806	B	11/2003
WO	WO-8300288	A1	2/1983
WO	WO-8806599	A1	9/1988
WO	WO-8910937	A1	11/1989
WO	WO-9007522	A1	7/1990
WO	WO-9011299	A1	10/1990
WO	WO-9103550	A1	3/1991
WO	WO-9116929	A1	11/1991
WO	92/00321		1/1992
WO	92/00321	A1	1/1992
WO	WO-9212999	A1	8/1992
WO	WO-9318786	A1	9/1993
WO	WO-9414461	A1	7/1994
WO	WO-9500550	A1	1/1995
WO	WO-9524183	A1	9/1995
WO	WO-9604307	A1	2/1996
WO	WO-9607399	A1	3/1996
WO	WO-9611705	A1	4/1996
WO	96/32414	A1	10/1996
WO	WO-9634882	A1	11/1996
WO	WO-9641606	A2	12/1996
WO	WO-9701331	A2	1/1997
WO	WO-9748413	A1	12/1997
WO	98/05351	A1	2/1998
WO	98/08871	A1	3/1998
WO	WO-9808531	A1	3/1998
WO	WO-9808873	A1	3/1998
WO	WO-9819698	A1	5/1998
WO	98/30231	A1	7/1998
WO	WO-9835033	A1	8/1998

US 9,526,764 B2

(56)	References Cited			WO	2006/029634	A2	3/2006
	FOREIGN PATENT DOCUMENTS			WO	2006/051103	A2	5/2006
				WO	WO-2006051110	A2	5/2006
				WO	WO-2006058620	A2	6/2006
WO	WO-9839022	A1	9/1998	WO	WO-2006110551	A2	10/2006
WO	WO-9842749	A1	10/1998	WO	WO-2007001150	A2	1/2007
WO	WO-9856406	A1	12/1998	WO	WO-2007024700	A2	3/2007
WO	WO-9856418	A1	12/1998	WO	WO-2007028394	A2	3/2007
WO	99/07404		2/1999	WO	WO-2007031187	A1	3/2007
WO	99/07404	A1	2/1999	WO	WO-2007035665	A1	3/2007
WO	99/25727	A2	5/1999	WO	WO-2007036299	A2	4/2007
WO	99/25728	A1	5/1999	WO	WO-2007037607	A1	4/2007
WO	WO-9921573	A1	5/1999	WO	WO-2007044867	A2	4/2007
WO	WO-9921578	A1	5/1999	WO	WO-2007050656	A2	5/2007
WO	WO-9924071	A1	5/1999	WO	WO-2007075534	A2	7/2007
WO	WO-9940788	A1	8/1999	WO	WO-2007081824	A2	7/2007
WO	9943708	A1	9/1999	WO	WO-2007082381	A1	7/2007
WO	WO-9946283	A1	9/1999	WO	WO-2007095288	A2	8/2007
WO	WO-9962558	A1	12/1999	WO	WO-2007104786	A1	9/2007
WO	WO-0023098	A1	4/2000	WO	WO-2007109221	A2	9/2007
WO	WO-0023099	A1	4/2000	WO	WO-2007113205	A1	10/2007
WO	WO-0029013	A1	5/2000	WO	WO-2007120899	A2	10/2007
WO	WO-0041546	A2	7/2000	WO	WO-2008006496	A1	1/2008
WO	WO-0066629	A1	11/2000	WO	WO-2008013938	A2	1/2008
WO	WO-0074736	A1	12/2000	WO	WO 2008/021560		2/2008
WO	01/02039	A1	1/2001	WO	WO-2008023050	A1	2/2008
WO	01/04156	A1	1/2001	WO	WO-2008028914	A1	3/2008
WO	WO-0100223	A2	1/2001	WO	WO-2008034881	A1	3/2008
WO	WO-0112155	A1	2/2001	WO	WO-2008124522	A2	10/2008
WO	WO-0121154	A2	3/2001	WO	WO-2008133908	A2	11/2008
WO	WO-0125278	A1	4/2001	WO	WO-2008145323	A1	12/2008
WO	WO-0128555	A1	4/2001	WO	WO-2009004627	A2	1/2009
WO	WO-0137808	A1	5/2001	WO	WO-2009030498	A2	3/2009
WO	WO-0143762	A2	6/2001	WO	WO-2009030499	A1	3/2009
WO	WO-0151071	A2	7/2001	WO	2009/039963	A1	4/2009
WO	WO-0152937	A1	7/2001	WO	WO-2009048959	A1	4/2009
WO	WO-0193837	A2	12/2001	WO	2009/063072	A2	5/2009
WO	WO-0200243	A2	1/2002	WO	WO-2009056569	A1	5/2009
WO	WO-0224214	A2	3/2002	WO	2009/087081	A2	7/2009
WO	WO-02064115	A1	8/2002	WO	WO 2009/087082		7/2009
WO	WO-02065985	A2	8/2002	WO	WO-2009089181	A1	7/2009
WO	WO-02066628	A2	8/2002	WO	WO-2009098318	A1	8/2009
WO	WO-02068660	A1	9/2002	WO	WO-2009102467	A2	8/2009
WO	WO-02070722	A1	9/2002	WO	WO 2009/143014		11/2009
WO	WO-02076495	A1	10/2002	WO	WO-2009134380	A2	11/2009
WO	WO-02079250	A1	10/2002	WO	WO-2010030670	A2	3/2010
WO	WO-03002021	A2	1/2003	WO	WO-2010043566	A2	4/2010
WO	03/020201	A2	3/2003	WO	WO-2010044867	A1	4/2010
WO	WO-03035028	A1	5/2003	WO	WO-2010092163	A2	8/2010
WO	WO-03035051	A2	5/2003	WO	WO-2011012719	A1	2/2011
WO	WO-03044210	A2	5/2003	WO	WO-2011017554	A2	2/2011
WO	WO-03053339	A2	7/2003	WO	WO-2011029892	A2	3/2011
WO	WO-03066084	A1	8/2003	WO	WO-2011058082	A1	5/2011
WO	WO-03094951	A1	11/2003	WO	WO-2011058083	A1	5/2011
WO	WO-03094956	A1	11/2003	WO	WO-2011103575	A1	8/2011
WO	WO-03101395	A2	12/2003	WO	WO-2011122921	A2	10/2011
WO	WO-03105888	A1	12/2003	WO	WO-2011128374	A1	10/2011
WO	2004/003523	A1	1/2004	WO	WO-2011144673	A2	11/2011
WO	2004/005342	A1	1/2004	WO	WO-2011144674	A2	11/2011
WO	2004/035623	A2	4/2004	WO	WO-2011147980	A1	12/2011
WO	WO-2004045592	A2	6/2004	WO	WO-2011157402	A1	12/2011
WO	WO-2004064862	A1	8/2004	WO	WO-2011160066	A1	12/2011
WO	WO-2004078196	A1	9/2004	WO	WO-2012012352	A2	1/2012
WO	WO-2004078197	A1	9/2004	WO	WO-2012028172	A1	3/2012
WO	WO-2004078198	A1	9/2004	WO	WO-2012055967	A2	5/2012
WO	WO-2004080480	A1	9/2004	WO	WO-2012065996	A1	5/2012
WO	WO-2004096854	A2	11/2004	WO	WO-2012066086	A1	5/2012
WO	WO-2004105781	A2	12/2004	WO	WO-2012080320	A1	6/2012
WO	WO-2004107979	A1	12/2004	WO	WO-2012104342	A1	8/2012
WO	2005/023291	A2	3/2005	WO	WO-2012125569	A2	9/2012
WO	WO-2005021022	A2	3/2005	WO	WO-2012156296	A1	11/2012
WO	WO-2005028516	A2	3/2005	WO	WO-2012156299	A1	11/2012
WO	2005/046716	A1	5/2005	WO	WO-2012177929	A2	12/2012
WO	WO 2005/046716	*	5/2005	WO	WO-2013060850	A1	5/2013
		 A61K 38/26	WO	WO-2014017849	A1	1/2014
WO	WO-2005048950	A2	6/2005	WO	WO-2014118355	A1	8/2014
WO	WO-2005112949	A1	12/2005	WO	WO 2014/202483		12/2014
WO	WO-2005117948	A1	12/2005				
WO	2006/000567	A2	1/2006				
WO	WO-2006015879	A1	2/2006				

US 9,526,764 B2

Page 5

(56)

References Cited**OTHER PUBLICATIONS**

Tews, D et al., “Enhanced protection against cytokine-and fatty acid-induced apoptosis in pancreatic beta cells by combined treatment with glucagon-like peptide-1 receptor agonists and insulin analogues,” *Hormone and Metabolic Research* (2008), vol. 40, No. 3, pp. 172-180, Database Biosis Abstract XP-002577467, Database Accession No. PREV200800343949.

International Search Report dated May 6, 2010.

International Preliminary Report on Patentability dated May 24, 2011 issued in PCT/EP2009/063195.

Office Action dated May 30, 2011 issued in corresponding New Zealand Patent Application No. 592283.

Office Action dated Jul. 27, 2010 issued in corresponding Panama Patent Application No. PI/2009/88459-01.

Representative Report dated Oct. 1, 2010 on an issued Office Action issued in corresponding Patent Application No. 933/2009 in Pakistan.

Tews et al., “Enhanced Protection” *Diabetes* (2007) pp. A72-A73, vol. 56, suppl. 1.

www.fda.gov/CombinationProducts/AboutCombinationProducts/usm101496.1/htm, 2009 downloaded Jul. 13, 2012.

Opposition filed on Aug. 14, 2012 against Application No. SP-0321-2009 in Bolivia.

Correa, C., “Pautas para el examen de patentes farmaceuticas. Una perspectiva desde la Salud Publica. Documento de Trabajo” Universidad de Buenos Aires, Mar. 2008, pp. 19-20.

Schubert-Zsilavecz, Manfred et al., “Bessere Blutzuckerwerte beim Diabetiker—Insulin glargin—ein langwirksames Insulinanalogon,” *Pharmazie in unserer Zeit* (2001), vol. 2, pp. 125-130.

Holst, Jens Juul, “Glucagon-like Peptide-1, A Gastrointestinal Hormone with a Pharmaceutical Potential,” *Current Medicinal Chemistry* (1999), vol. 6, pp. 1005-1017.

Nauck, M. A. et al., “Glucagon-like peptide 1 (GLP-1) as a new therapeutic approach for Type 2-diabetes,” *Experimental and Clinical Endocrinology and Diabetes* (1997), vol. 105, pp. 187-195.

Lopez-Delgado, Maria I. et al., “Effects of Glucagon-Like Peptide 1 on the Kinetics of Glycogen Synthase a in Hepatocytes from Normal and Diabetic Rats,” *Endocrinology* (1998), vol. 139, No. 6, pp. 2811-2817.

Tews, D. et al., “Enhanced Protection against Cytokine- and Fatty Acid-induced Apoptosis in Pancreatic Beta Cells by Combined Treatment with Glucagon-like Peptide-1 Receptor Agonists and Insulin Analogues,” *Hormone and Metabolic Research* (2008), vol. 40, No. 3, pp. 172-180.

International Preliminary Report on Patentability dated Sep. 23, 2010 issued in PCT/EP2009/063195.

International Search Report dated May 6, 2010 issued in PCT/EP2009/063195.

Opposition filed by Association of the National Pharmaceutical Industry (ASIFAN) on Oct. 21, 2011.

English Translation of Opposition filed by Association de Laboratorios Farmaceuticos (ALAFAR) on Oct. 29, 2011.

The Diabetes Control and Complications Trial Research Group, “The Effect of Intensive Treatment of Diabetes on the Development and Progression of Long-Term Complications in Insulin-Dependent Diabetes Mellitus” 329(14):977-86 (1993).

Goke et al., “Exendin-4 is a high potency agonist and truncated exendin-(9-39)-amide an antagonist at the glucagon-like peptide 1-(7-36)-amide receptor of insulin-secreting beta-cells.” *J. Biol Chem* 268(26):19650-55 (1993).

Substantive Examination Report from the Intellectual Property Office of the Philippines, Bureau of Patents for Philippines Application No. 1-2011-500747, mailed Feb. 28, 2014, pp. 1-3.

Official Action for Russian Application No. 2011119639 of Feb. 14, 2014, pp. 1-4.

Heinrich et al., “Pre-proglucagon messenger ribonucleic acid: nucleotide and encoded amino acid sequences of the rat pancreatic complementary deoxyribonucleic acid.” *Endocrinology* 115(6):2176-81 (1984).

Raufman, “Bioactive peptides from lizard venoms.” *Regulatory Peptides* 61(1):1-18 (1996).

Uttenthal et al., “Molecular forms of glucagon-like peptide-1 in human pancreas and glucagonomas.” *J Clin Endocrinol Metabol* 61(3):472-79 (1985).

Office Action dated Oct. 9, 2013 issued in corresponding Chinese Patent Application No. CN200980150799.X.

Office Action dated Sep. 30, 2013 issued in corresponding Malaysian Patent Appln. No. PI 2011001321.

Pi-Sunyer et al., “The effects of pharmacologic agents for type 2 diabetes mellitus on body weight”, *Postgrad Med.*, Jul. 2008, 5-17, 120(2).

Weyer et al., “Long-term changes in insulin action and insulin secretion associated with gain, loss, regain and maintenance of body weight”, *Diabetologia*, Jan. 2000, 36-46, (43)1.

Russian Office Action dated Oct. 17, 2013 issued in Russian Application No. 2011119636/15 together with English translation.

Office Action dated Feb. 26, 2013 issued in corresponding Ukrainian Patent Application No. UA 2011 06137.

Office Action dated Apr. 14, 2013 issued in corresponding Israeli Patent Application No. IL 212 258.

Office Action dated Apr. 18, 2013 issued in corresponding Chinese Patent Application No. CN 200980150799.X.

Office Action dated Apr. 25, 2013 issued in corresponding Indonesian Patent Application No. ID W-00201101325.

Opposition filed May 3, 2013 in corresponding Chilean Patent Application No. CL 851-2011.

Office Action dated May 20, 2013 issued in corresponding Australian Patent Application No. AU 2009305472.

Office Action dated May 29, 2013 issued in corresponding El Salvador Patent Application based on PCT/EP2009/063195.

Office Action dated Jun. 12, 2013 issued in corresponding Dominican Republic Patent Application No. DO 2011-0104.

Opposition filed Jun. 20, 2013 in corresponding Guatemala Patent Application No. GT A2011-0096.

Office Action dated Aug. 8, 2013 issued in corresponding Colombian Patent Application No. CO 2011-0104.

Office Action dated Aug. 9, 2013 issued in corresponding Ukrainian Patent Application No. UA 2011-06137.

English translation of Japanese Notification of Reasons for Refusal dated Dec. 24, 2013 issued in Japanese Application No. 2011-531457.

English translation of Japanese Notification of Reasons for Refusal dated Dec. 24, 2013 issued in Japanese Application No. 2012-221012.

English translation of Decision of Refusal for Japanese Patent Application No. 2012-221012, dispatched Aug. 19, 2014, pp. 1-4. Patent Examination Report No. 2 for Australian Patent Application No. 2009305472, issued Sep. 30, 2013, pp. 1-4.

NHSC—National Horizon Scanning Center, “AVE0010 (ZP10) for type 2 diabetes mellitus” University of Birmingham, England; pp. 1-6 (Dec. 2008).

Search Report of the Intellectual Property Office of Singapore for Patent Application No. 10201500871T, dated Nov. 2, 2015, pp. 1-3.

Chi et al., “Excipients and their Effects on the Quality of Biologics” pp. 1-9, (May 2012).

Non-Final Rejection issued in U.S. Appl. No. 13/382,442; mailed Mar. 31, 2016, pp. 1-29.

International Search Report by the ISA for International Application No. PCT/EP2015/079285; dated Mar. 9, 2016, pp. 1-7.

English translation of the TIPO Search Report for ROC Patent Application No. 104116749, dated Feb. 22, 2016, one page.

U.S. Appl. No. 13/382,772, filed May 29, 2012, Schoettle.

U.S. Appl. No. 13/382,442, filed Feb. 21, 2012, Schoettle.

U.S. Appl. No. 13/509,507, filed Jul. 30, 2014, Brunner-Schwarz et al.

U.S. Appl. No. 13/509,542, filed Aug. 2, 2012, Hagendorf et al.

U.S. Appl. No. 14/172,151, filed Feb. 4, 2014, Bley et al.

U.S. Appl. No. 12/617,805, filed Nov. 13, 2009, Silvestre et al.

U.S. Appl. No. 12/617,811, filed Nov. 13, 2009, Silvestre et al.

U.S. Appl. No. 14/220,562, filed Mar. 20, 2014, Becker et al.

U.S. Appl. No. 13/700,631, filed Nov. 11, 2012, Becker et al.

U.S. Appl. No. 13/819,114, filed Apr. 29, 2013, Boka et al.

US 9,526,764 B2

Page 6

(56)

References Cited

OTHER PUBLICATIONS

U.S. Appl. No. 13/363,956, filed Feb. 1, 2012, Silvestre et al.
 U.S. Appl. No. 13/432,811, filed Mar. 28, 2012, Boka et al.
 U.S. Appl. No. 13/469,633, filed May 11, 2012, Ruus et al.
 U.S. Appl. No. 13/467,707, filed May 9, 2012, Niemoller et al.
 U.S. Appl. No. 13/468,422, filed May 10, 2012, Silvestre et al.
 U.S. Appl. No. 13/467,757, filed May 9, 2012, Silvestre et al.
 U.S. Appl. No. 13/595,590, filed Aug. 27, 2012, Niemoller et al.
 U.S. Appl. No. 13/602,913, filed Sep. 4, 2012, Hess et al.
 U.S. Appl. No. 13/633,563, filed Oct. 2, 2012, Stechl et al.
 U.S. Appl. No. 13/633,496, filed Oct. 2, 2012, Stechl et al.
 U.S. Appl. No. 13/661,476, filed Oct. 26, 2012, Silvestre et al.
 U.S. Appl. No. 14/303,895, filed Jun. 13, 2014, Souhami et al.
 English Translation of TIPO Search Report for ROC Patent Application No. 101130936, dated Dec. 1, 2015, one page.
 Final Rejection issued in U.S. Appl. No. 13/467,707; mailed Feb. 12, 2016, pp. 1-12.
 Final-Rejection issued in U.S. Appl. No. 13/432,811; mailed Feb. 10, 2016, pp. 1-9.
 Final Rejection issued in U.S. Appl. No. 13/509,542, mailed Feb. 10, 2016, pp. 1-40.
 18th World Health Congress (Helsinki). WMA Declaration of Helsinki—Ethical Principles for Medical Research Involving Human Subjects; WMA; Jun. 1964, pp. 1-8.
 Abbas T., et al., “Impairment of Synaptic Plasticity and Memory formation in GLP-1 Receptor Ko Mice: Interaction Between Type 2 Diabetes and Alzheimer’s Disease,” Behavioural Brain Research, 2009, vol. 205 (1), pp. 265-271.
 Action to Control Cardiovascular Risk in Diabetes Study Group, “Effects of Intensive Glucose Lowering in Type 2 Diabetes,” The New England Journal of Medicine, 2008, vol. 358 (24), pp. 2545-2559.
 Actrapid® prescribing information, Apr. 2011, pp. 1-4.
 Actrapid® summary of product characteristics, Apr. 2011, pp. 1-11.
 Aderinwale O.G., et al., “Current Therapies and New Strategies for the Management of Alzheimer’s Disease,” American Journal of Alzheimer’s Disease and Other Dementias, 2010, vol. 25 (5), pp. 414-424.
 Agholme L., et al., “An in Vitro Model for Neuroscience: Differentiation of SH-SY5Y Cells into Cells with Morphological and Biochemical Characteristics of Mature Neurons,” Journal of Alzheimer’s Disease, 2010, vol. 20, pp. 1069-1082.
 Ahren et al., Abstract “Efficacy and Safety of Lixisenatide QD Morning and Evening Injections vs Placebo in T2DM Inadequately Controlled on Metformin (GetGoal-M)” Oral presentation O-0591 presented at the World Diabetes Congress in Dubai, Dec. 5-8, 2011, one page.
 Ahualli J., “The Double Duct Sign,” Radiology, 2007, vol. 244 (1), pp. 314-315.
 Akbar D.H., “Sub-Optimal Postprandial Blood Glucose Level in Diabetics Attending the Outpatient Clinic of a University Hospital,” Saudi Med Journal, 2003, vol. 24 (10), pp. 1109-1112.
 American Diabetes Association (ADA) Committee Report—The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus—Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus, Diabetes Care, 21(Supplement 1): S5-S19 (Jan. 1998).
 Aoki K., et al., “Hydrolysis of Nonionic Surfactants,” Annual Report Takeda Research Laboratory, 1968, vol. 27, pp. 172-176.
 Apidra® prescribing information, Apr. 2012, pp. 1-6.
 Arnolds et al., “Insulin Glargine (GLAR) plus Metformin (MET): An Efficacious and Safe Regimen that can be combined with Exenatide (EXE) or Sitagliptin (SITA)” Diabetes, 58(Suppl. 1): A141, Jun. 2009.
 Arnolds S., et al., “Basal Insulin Glargine Vs Prandial Insulin Lispro in Type 2 Diabetes,” Lancet, 2008, vol. 378 (9636), pp. 370-371.
 Arnolds S., et al., “Further Improvement in Postprandial Glucose Control with Addition of Exenatide or Sitagliptin to Combination therapy with Insulin Glargine and Metformin—A Proof-of-Concept Study,” Diabetes Care, 2010, vol. 33 (7), pp. 1509-1515.

Auerbach R., et al., “Angiogenesis Assays: Problems and Pitfalls,” Cancer Metastasis Reviews, 2000, vol. 19 (1-2), pp. 167-172.
 Bakaysa D.L., et al., “Physicochemical Basis for the Rapid Time-Action of Lysb28 Prob29-Insulin: Dissociation of a Protein-Ligand Complex,” Protein science, 1996, vol. 5 (12), pp. 2521- 2531.
 Banks W.A., et al., “Brain Uptake of the Glucagon-Like Peptide-1 Antagonist Exendin(9-39) After intranasal Administration,” The Journal of Pharmacology and Experimental Therapeutics, 2004, vol. 309 (2), pp. 469-475.
 Barnett A., “Dosing of Insulin Glargine in the Treatment of Type 2 Diabetes,” Clinical Therapeutics, 2007, vol. 29 (6), pp. 987-999.
 Barnett A.H., et al., “Tolerability and Efficacy of Exenatide and Titrated Insulin Glargine in Adult Patients with Type 2 Diabetes Previously Uncontrolled with Metformin or a Sulfonylurea: A Multinational, Randomized, Open-Label, Two-Period, Crossover Noninferiority Trial,” Clinical Therapeutics, 2007, vol. 29 (11), pp. 2333-2348.
 Barnett A.H., “Insulin Glargine in the Treatment of Type 1 and Type 2 Diabetes,” Vascular Health and Risk Management, 2006, vol. 2 (1), pp. 59-67.
 Barnett A.H., “Lixisenatide: Evidence for its Potential Use in the Treatment of Type 2 Diabetes,” Core Evidence, 2011, vol. 6, pp. 67-79.
 Barnett R.O., et al., “Insulin Analogues,” Lancet, 1997, vol. 349 (9044), pp. 47-51.
 Behar J., et al., “Functional Gallbladder and Sphincter of Oddi Disorders,” Gastroenterology, 2006, vol. 130 (5), pp. 1498-1509.
 Beintema J.J., et al., “Molecular Evolution of Rodent Insulins,” Molecular Biology and Evolution, 1987, vol. 4 (1), pp. 10-18.
 Berger M., “Towards More Physiological Insulin Therapy in the 1990s a Comment,” Diabetes Research and Clinical Practice, 1989, vol. 6 (4), pp. S25-S31.
 Berlie H., et al., “Glucagon-Like Peptide-1 Receptor Agonists as Add-On therapy to Basal Insulin in Patients with Type 2 Diabetes: A Systematic Review,” Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2012, vol. 5, pp. 165-174.
 Berlinsulin® H prescribing information, Apr. 2012, pp. 1-4.
 Berlinsulin® H summary of product characteristics, Apr. 2012, pp. 1-11.
 Bertram L., et al., “The Genetics of Alzheimer Disease: Back to the Future,” Neuron, 2010, vol. 68 (2), pp. 270-281.
 Best, Mathematics and Statistics pp. 1-39, 1988.
 Bethel M.A., et al., “Basal Insulin Therapy in Type 2 Diabetes,” The Journal of the American Board of the Family Practice, 2005, vol. 18 (3), pp. 199-204.
 Bhatt N.P., et al., “Chemical Pathways of Peptide Degradation. I. Deamidation of Adrenocorticotrophic Hormone,” Pharmaceutical Research, 1990, vol. 7 (6), pp. 593-599.
 Blanchard V., et al., “Time Sequence of Maturation of Dystrophic Neurites Associated with Abeta Deposits in APP/PS1 Transgenic Mice,” Experimental Neurology, 2003, vol. 184, pp. 247-263.
 Bland J.M., et al., “Measurement Error,” British Medical Journal, 1996, vol. 312 (7047), pp. 1654.
 Bolen S., et al., “Systematic Review: Comparative Effectiveness and Safety of oral Medications for Type 2 Diabetes Mellitus,” Annals of Internal Medicine, 2007, vol. 147 (6), pp. 386-399.
 Bolli et al., “Efficacy and safety of lixisenatide once-daily versus placebo in patients with type 2 diabetes mellitus insufficiently controlled on metformin (GetGoal-F1).” Presentation Abstract No. 784, EASD Meeting Sep. 12-16, 2014.
 Bolli G.B., et al., “Efficacy and Safety of Lixisenatide once Daily Vs Placebo in People with Type 2 Diabetes Insufficiently Controlled on Metformin (Getgoal-F1),” Diabetic Medicine, 2014, vol. 31 (2), pp. 176-184.
 Bolli G.B., “The Pharmacokinetic Basis of Insulin Therapy in Diabetes Mellitus,” Diabetes Research and Clinical Practice, 1989, vol. 6 (4), pp. S3-15.
 Boutajangout A., et al., “Characterisation of Cytoskeletal Abnormalities in Mice Transgenic for Wild-Type Human Tau and Familial Alzheimer’s Disease Mutants of APP and Presenilin-1,” Neurobiology of Disease, 2004, vol. 15 (1), pp. 47-60.
 Boutajangout A., et al., “Increased Tau Phosphorylation But Absence of formation of Neurofibrillary Tangles in Mice Double

US 9,526,764 B2

Page 7

(56)

References Cited

OTHER PUBLICATIONS

Transgenic for Human Tau and Alzheimer Mutant (M146L) Presenilin-1," *Neuroscience Letters*, 2002, vol. 318 (1), pp. 29-33.

Brange "Galenics of Insulin" 1987, p. 35-36.

Brange J., et al., "Chemical Stability of Insulin 3. Influence of Excipients, formulation, and Ph," *Acta Pharmaceutica Nordica*, 1992, vol. 4 (3), pp. 149-158.

Brange J., et al., "Design of Insulin Analogues for Meal-Related therapy," *Journal of Diabetes and Its Complications*, 1993, vol. 7 (2), pp. 106-112.

Brange J., et al., "Insulin Structure and Stability," *Pharmaceutical Biotechnology*, 1993, vol. 5, pp. 315-350.

Brange J., et al., "Monomeric Insulins and their Experimental and Clinical Implications," *Diabetes Care*, 1990, vol. 13 (9), pp. 923-954.

Brange J., et al., "Neutral Insulin Solutions Physically Stabilized by Addition of Zn²⁺," *Diabetic Medicine*, 1986, vol. 3, pp. 532-536.

Brange J., et al., "Toward Understanding Insulin Fibrillation," *Journal of Pharmaceutical Sciences*, 1997, vol. 86 (5), pp. 517-525.

Brod M., et al., "Adherence Patterns in Patients with Type 2 Diabetes on Basal Insulin Analogues: Missed, Mistimed and Reduced Doses," *Current Medical Research and Opinion*, 2012, vol. 28 (12), pp. 1933-1946.

Brod M., et al., "Examining Correlates of Treatment Satisfaction for injectable Insulin in Type 2 Diabetes: Lessons Learned from a Clinical Trial Comparing Biphasic and Basal Analogues," *Health Quality of Life Outcomes*, 2007, vol. 5, pp. 1-10.

Broderick J., et al., "Guidelines for the Management of Spontaneous intracerebral Hemorrhage in Adults," *Circulation*, 2007, vol. 116 (16), pp. e391-e413.

Brown J.B., et al., "Slow Response to Loss of Glycemic Control in Type 2 Diabetes Mellitus," *American Journal of Managed Care*, 2003, vol. 9 (3), pp. 213-217.

"Buffer" *Oxford Dictionary of Biochemistry and Molecular Biology*, Oxford University Press, 2001, p. 83.

Burgermeister W., et al., "The Isolation of Insulin from the Pancreas," *Insulin*, 1975, vol. Part 2, pp. 715-727.

Burke G.T., et al., "Nature of the B10 Amino Acid Residue Requirements for High Biological Activity of Insulin," *International Journal of Peptide and Protein Research*, 1984, vol. 23 (4), pp. 394-401.

Buse J.B., et al., "Use of Twice-Daily Exenatide in Basal Insulin-Treated Patients with Type 2 Diabetes: A Randomized, Controlled Trial," *Annals of Internal Medicine*, 2011, vol. 154 (2), pp. 103-112.

Byetta—Summary of Product Characteristics, updated Jan. 27, 2015, last accessed Apr. 18, 2015, pp. 1-12.

Byrne M.M., et al., "Inhibitory Effects of Hyperglycaemia on Fed Jejunal Motility: Potential Role of Hyperinsulinaemia," *European Journal of Clinical Investigation*, 1998, vol. 28 (1), pp. 72-78.

Cadario B., "SITAGLIPTIN," *Drug Information Perspectives*, 2010, vol. 30 (4), pp. 1-6.

Campas C., et al., "Ave-0010 GLP-1 Receptor Agonist Treatment of Diabetes," *Drugs of the Future*, 2008, vol. 33 (10), pp. 838-840.

Campbell R.K., et al., "Insulin Glargine," *Clinical Therapeutics*, 2001, vol. 23 (12), pp. 1938-1957.

Casas C., et al., "Massive CA1/2 Neuronal Loss with Intraneuronal and N-Terminal Truncated Abeta42 Accumulation in a Novel Alzheimer Transgenic Model," *American Journal of Pathology*, 2004, vol. 165 (4), pp. 1289-1300.

Chancel, "Natixis Conference on Diabetes." Sanofi, Paris, pp. 1-23 (Nov. 8, 2011).

Charles M.A., et al., "Prevention of Type 2 Diabetes: Role of Metformin," *Drugs*, 1999, vol. 58 (Suppl 1), pp. 71-73.

Chatterjee S., et al., "Insulin Glargine and its Place in the Treatment of Types 1 and 2 Diabetes Mellitus," *Expert Opinion on Pharmacotherapy*, 2006, vol. 7 (10), pp. 1357-1371.

Chen Y.E., et al., "Tissue-Specific Expression of Unique mRNAs That Encode Proglucagon-Derived Peptides or Exendin 4 in the Lizard," *The Journal of Biological Chemistry*, 1997, vol. 272 (7), pp. 4108-4115.

Cheung Y.T., et al., "Effects of All-Trans-Retinoic Acid on Human SH-SY5Y Neuroblastoma as in Vitro Model in Neurotoxicity Research," *Neurotoxicology*, 2009, vol. 30 (1), pp. 127-135.

Childs B.P., et al., "Defining and Reporting Hypoglycemia in Diabetes: A Report from the American Diabetes Association Workgroup on Hypoglycemia," *Diabetes Care*, 2005, vol. 28 (5), pp. 1245-1249.

Cholangiocarcinoma, Johns Hopkins Medicine Webstite, https://gi.jhsps.org/GDLDisease.aspx?CurrentUDV=31&GDLCat_ID=AF793A59-B736-42CB-9E1FE79D2B9FC358&GDL_Disease_ID=A6D10E80-887D-49A7-B3BB-0517D38CE757, accessed on May 14, 2014, pp. 1-12.

Christensen M., et al., "Lixisenatide, A Novel GLP-1 Receptor Agonist for the Treatment of Type 2 Diabetes Mellitus," *IDrugs: The Investigational Drugs Journal*, 2009, vol. 12 (8), pp. 503-513.

Christensen M., et al., "Lixisenatide for Type 2 Diabetes Mellitus," *Expert Opinion on Investigational Drugs*, 2011, vol. 20 (4), pp. 549-557.

Cochran E., et al., "The Use of U-500 in Patients with Extreme Insulin Resistance," *Diabetes Care*, 2005, vol. 28 (5), pp. 1240-1244.

Colclough et al., Abstract "Levels of FPG and HbA1c Control and the Relationship to BMI in T2D Patients Treated with Basal Insulin and OAD Therapy." Abstract 2416-PO; Presented at the 72nd Scientific Session at the American Diabetes Association Meeting, 2012, A609, one page.

Colino E., et al., "Therapy with Insulin Glargine (Lantus) in toddlers, Children and Adolescents with Type 1 Diabetes," *Diabetes Research and Clinical Practice*, 2005, vol. 70 (1), pp. 1-7.

Community register of medicinal products for human use, Chemical Subgroup A10BX, "Lyxumia" European Commision—Public Health, p. 1-2 (May 2, 2013).

Craig et al., "ISPAD Clinical Practice Consensus Guidelines 2014 Compendium—Definition, epidemiology, and classification of diabetes in children and adolescents." *Pediatric Diabetes*, 15(Suppl. 20):4-17 (2014).

Crapo P.A., et al., "Postprandial Plasma-Glucose and -Insulin Responses to Different Complex Carbohydrates," *Diabetes*, 1977, vol. 26 (12), pp. 1178-1183.

Croom K.F., et al., "Liraglutide a Review of its Use in Type 2 Diabetes Mellitus," *Drugs*, 2009, vol. 69 (14), pp. 1985-2004.

Cryer P.E., "Hypoglycemia Is the Limiting Factor in the Management of Diabetes," *Diabetes/Metabolism Research and Reviews*, 1999, vol. 15 (1), pp. 42-46.

Cvetkovic R.S., et al., "Exenatide A Review of its Use in Patients with Type 2 Diabetes Mellitus (As an Adjunct to Metformin and/or a Sulfonylurea)," *Drugs*, 2007, vol. 67 (6), pp. 935-954.

Czech C., et al., "Proteolytical Processing of Mutated Human Amyloid Precursor Protein in Transgenic Mice," *Brain Research Molecular Brain Research*, 1997, vol. 47 (1-2), pp. 108-116.

D'Alessio D., "GLP-1 Receptor Agonists: Strategies for PPG Control," *Medical Nursing Education*, 2011, vol. 3, pp. 1-26.

D'Alessio D.A., et al., "Glucagon-Like Peptide 1 Enhances Glucose tolerance both by Stimulation of Insulin Release and by increasing Insulin-Independent Glucose Disposal," *Journal of Clinical Investigation*, 1994, vol. 93 (5), pp. 2263-2266.

Database, ADISCTI, "A randomized, 4-sequence, cross-over, double bind, dose response study of 0.4, 0.6 and 0.09 U/kg insulin glargine U300 compared to 0.4 U/kg Lantus U100 in patients with diabetes mellitus type I using euglycemic clamp technique" last updated Dec. 16, 2010, pp. 1-4.

Davis How to Convert mg to mmol/L, available online at http://www.ehow.com/how_8498850_convert_mg-mmol.html (accessed on Nov. 11, 2015).

De Arriba S.G., et al., "Carbonyl Stress and Nmda Receptor Activation Contribute to Methylglyoxal Neurotoxicity," *Free Radical Biology and Medicine*, 2006, vol. 40 (5), pp. 779-790.

De La Pena A., et al., "Pharmacokinetics and Pharmacodynamics of High-Dose Human Regular U-500 Insulin Versus Human Regular U-100 Insulin in Healthy Obese Subjects," *Diabetes Care*, 2011, vol. 34 (12), pp. 2496-2501.

De Rosa R., et al., "Intranasal Administration of Nerve Growth Factor (Ngf) Rescues Recognition Memory Deficits in Ad11 Anti-

US 9,526,764 B2

Page 8

(56)

References Cited

OTHER PUBLICATIONS

Ngf Transgenic Mice,” Proceedings of the National Academy of Sciences of the United States of America, 2005, vol. 102 (10), pp. 3811-3816.

Deacon C.F., et al., “Dipeptidyl Peptidase IV inhibition Potentiates the Insulinotropic Effect of Glucagon-Like Peptide 1 in the Anesthetized Pig,” Diabetes, 1998, vol. 47 (5), pp. 764-769.

Deacon C.F., et al., “Dipeptidyl Peptidase IV Resistant Analogues of Glucagon-Like Peptide-1 Which have Extended Metabolic Stability and Improved Biological Activity,” Diabetologia, 1998, vol. 41 (3), pp. 271-278.

Definition of indication, Merriam-Webster online, accessed Oct. 22, 2015, 2 pages.

Definition of palliative, <http://medicaldictionary.thefreedictionary.com/>, accessed on Nov. 6, 2014, pp. 1-2.

Definition of sphincter of pancreatic duct in the Medical Dictionary, <http://medicaldictionary.thefreedictionary.com/>, accessed on May 22, 2014, pp. 1-2.

DeFronzo R.A., et al., “Effects of Exenatide (Exendin-4) on Glycemic Control and Weight Over 30 Weeks in Metformin-Treated Patients with Type 2 Diabetes,” Diabetes care, 2005, vol. 28 (5), pp. 1092-1100.

DeFronzo R.A., “Pathogenesis of Type 2 Diabetes: Implications for Metformin,” Drugs, 1999, vol. 58 (Suppl 1), pp. 29-30.

DeFronzo R.A., “Pharmacologic therapy for Type 2 Diabetes Mellitus,” Annals of Internal Medicine, 1999, vol. 131 (4), pp. 281-303.

Delatour B., et al., “Alzheimer Pathology Disorganizes Cortico-Cortical Circuitry: Direct Evidence from a Transgenic Animal Model,” Neurobiology of Disease, 2004, vol. 16 (1), pp. 41-47.

Devries J.H., et al., “Sequential intensification of Metformin Treatment in Type 2 Diabetes with Liraglutide Followed by Randomized Addition of Basal Insulin Prompted by A1C Targets,” Diabetes Care, 2012, vol. 35 (7), pp. 1446-1454.

Dewitt D.E., “Case Study: Treating New on-Set Catabolic Type 2 Diabetes with Glargine and Lispro,” Clinical Diabetes, 2006, vol. 24 (4), pp. 180-181.

Diabetes Prevention Program Research Group, “Reduction in the incidence of Type 2 Diabetes with Lifestyle intervention or Metformin,” The New England Journal of Medicine, 2002, vol. 346 (6), pp. 393-403.

Distiller et al., Poster: “Pharmacokinetics and Pharmacodynamics of a New GLP-1 Agonist AVE0010 in Type 2 Diabetes Patients” Meeting: 68th Scientific Sessions (Jun. 2008) Poster No. 520-P.

Dixon G.H., et al., “Regeneration of Insulin Activity from the Separated and inactive A and B Chains,” Nature, 1960, vol. 188 (4752), pp. 721-724.

Donelli G., et al., “Plastic Biliary Stent Occlusion: Factors Involved and Possible Preventive Approaches,” Clinical Medicine & Research, 2007, vol. 5 (1), pp. 53-60.

Dormandy J.A., et al., “Secondary Prevention of Macrovascular Events in Patients with Type 2 Diabetes in the Proactive Study (Prospective Pioglitazone Clinical Trial in Macrovascular Events): A Randomised Controlled Trial,” Lancet, 2005, vol. 366 (9493), pp. 1279-1289.

Doyle M.E. et al., “Mechanisms of Action of Glucagon-Like Peptide 1 in the Pancreas,” Pharmacology & Therapeutics, 2007, vol. 113 (3), pp. 546-593.

Drucker D.J. et al., “The incretin System: Glucagon-Like Peptide-1 Receptor Agonists and Dipeptidyl Peptidase-4 inhibitors in Type 2 Diabetes,” Lancet, 2006, vol. 368 (9548), pp. 1696-1705.

Drucker D.J., “Glucagon-Like Peptides,” Diabetes, 1998, vol. 47 (2), pp. 159-169.

Drucker D.J., “Mini Review: The Glucagon-Like Peptides,” Endocrinology, 2001, vol. 142 (2), pp. 521-527.

Drucker D.J., “The Biology of Incretin Hormones,” Cell Metabolism, 2006, vol. 3 (3), pp. 153-165.

DrugBank, “Insulin glargine,” available online at <http://www.drugbank.ca/drugs/DB00047>, 16 pages (accessed online Sep. 25, 2014).

Drury P.L., et al., “Diabetic Nephropathy,” British Medical Bulletin, 1989, vol. 45 (1), pp. 127-147.

Dubois B., et al., “Revising the Definition of Alzheimer’s Disease: A New Lexicon,” Lancet Neurology, 2010, vol. 9 (11), pp. 1118-1127.

Dunn C.J., et al., “Insulin Glargine: An Updated Review of its Use in the Management of Diabetes Mellitus,” Drugs, 2003, vol. 63 (16), pp. 1743-1778.

During M.J., et al., “Glucagon-Like Peptide-1 Receptor is involved in Learning and Neuroprotection,” Nature Medicine, 2003, vol. 9 (9), pp. 1173-1179.

Eckert A., et al., “Alzheimer’s Disease-Like Alterations in Peripheral Cells from Presenilin-1 Transgenic Mice,” Neurobiology of Disease, 2001, vol. 8 (2), pp. 331-342.

EFC10780 (Sanofi study), “A randomized, double-blind, double-dummy, 2-arm parallel-group, multicenter 24-week study comparing the efficacy and safety of AVE0010 to sitagliptin as add-on to metformin in obese type 2 diabetic patients younger than 50 and not adequately controlled with metformin (EFC10780)” p. 1-4 (Jan. 29, 2014).

EFC10781 Clinical Trials, “24-week Treatment With Lixisenatide in Type 2 Diabetes Insufficiently Controlled With Metformin and Insulin Glargine” ClinicalTrials.gov; EFC10781 pp. 1-5 (Sep. 2009).

EFC6017; Clinical Trial Eudra CT No. 2007-005884-92, accessed Apr. 24, 2015, one page.

EFC6018; Clinical trial EudraCT 2007-005887-29, “GETGOAL-MONO” accessed Jul. 27, 2014; pp. 1-16.

EMA—Science Medicines Health “TOUJEO” EPAR Summary for the Public, first published Nov. 5, 2009, pp. 1-3.

EMA Press Release, “European Medicines Agency recommends suspension of Avandia, Avandamet and Avaglim” pp. 1-2 (Sep. 23, 2010).

English translation of Search Report for Chinese Patent Application No. 201280053404.6; dated Feb. 10, 2015, pp. 1-3.

English translation of Search Report for Chinese Patent Application No. 20140220537.9; dated Feb. 13, 2015, pp. 1-2.

European Medicines Agency—Science Medicines Health, “Guideline on clinical investigation of medicinal products in the treatment of diabetes mellitus” Committee for Medicinal Products for Human Use, Jan. 20, 2010, pp. 1-19.

Ex Parte Herrmann, Appeal No. 2009-001777 U.S. Appl. No. 10/616,457, (B. PAI. Nov. 13, 2009).

Executive Summary, “Standards of Medical Care in Diabetes-2009” Diabetes Care, 32(Suppl. 1):S6-S12 (Jan. 2009).

Extended European Search Report for European Application No. 98 11 0889.7; dated Oct. 14, 1998, pp. 1-4.

Extended European Search Report for European Application No. 09 17 5876.3; dated Mar. 24, 2010, pp. 1-4.

Extended European Search Report for European Application No. 09 17 5877.1; dated Apr. 29, 2010, pp. 1-5.

Extended European Search Report for European Application No. 10 16 4368.2; dated Oct. 14, 2010, pp. 1-6.

Extended European Search Report for European Application No. 10 30 5780; dated Nov. 16, 2010, pp. 1-3.

Extended European Search Report for European Application No. 11 15 3106; dated Jul. 6, 2011, pp. 1-12.

Extended European Search Report for European Application No. 11 16 0270.2; dated Sep. 19, 2011, pp. 1-8.

Extended European Search Report for European Application No. 11 16 6415; dated Mar. 12, 2012, pp. 1-12.

Extended European Search Report for European Application No. 11 17 9149.7; dated Feb. 9, 2012, pp. 1-8.

Extended European Search Report for European Application No. 13 305 126; dated Apr. 11, 2013, pp. 1-7.

Extended European Search Report for European Application No. 13 305 432.0; dated Sep. 13, 2013, pp. 1-5.

Extended European Search Report for European Application No. 14 16 6877.2; of Aug. 18, 2014, pp. 1-6.

Extended European Search Report for European Application No. 14 19 7154.9; dated Apr. 8, 2015, pp. 1-7.

Fabunmi R., et al., “Patient Characteristics, Drug Adherence Patterns, and Hypoglycemia Costs for Patients with Type 2 Diabetes

US 9,526,764 B2

Page 9

(56)

References Cited

OTHER PUBLICATIONS

Mellitus Newly initiated on Exenatide or Insulin Glargine,” Current Medical Research and Opinion, 2009, vol. 25 (3), pp. 777-786.

Faivre E., et al., “Effects of Gip Analogues in Neuronal Signalling, Cell Proliferation and Learning and Memory,” Regulatory Peptides, 2010, vol. 164 (1), pp. 40-41.

FDA Guidance for Industry, US Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research (COER), pp. 1-11, Feb. 2014.

FDA label of Apidra®, May 2014, pp. 1-35.

FDA label of Humalog®, Mar. 2013, pp. 1-27.

FDA label of Lantus®, Oct. 2013, pp. 1-44.

Feinglos M.N., et al., “Effects of Liraglutide (Nn2211), A Long-Acting GLP-1 Analogue, on Glycaemic Control and Bodyweight in Subjects with Type 2 Diabetes,” Diabetic Medicine, 2005, vol. 22 (8), pp. 1016-1023.

Fieller E.C., “Symposium on Interval Estimation; Some Problems with Interval Estimation,” Journal of the Royal Statistical Society, 1954, vol. 16 (2), pp. 175-185.

Final Office Action from U.S. Appl. No. 12/617,805; mailed Feb. 11, 2013, pp. 1-13.

Final Office Action from U.S. Appl. No. 12/617,805; mailed Jan. 12, 2012, pp. 1-14.

Final Office Action from U.S. Appl. No. 12/617,805; mailed Jan. 13, 2015, pp. 1-11.

Final Office Action issued in U.S. Appl. No. 13/602,913; mailed on Apr. 2, 2015, pp. 1-7.

Final Office Action issued in U.S. Appl. No. 13/602,913; mailed on Jun. 20, 2014, pp. 1-27.

Final Office Action issued in U.S. Appl. No. 13/602,913; mailed on Sep. 13, 2013, pp. 1-11.

Final Rejection in U.S. Appl. No. 13/633,496; mailed Nov. 18, 2014, pp. 1-11.

Final Rejection in U.S. Appl. No. 13/633,496; mailed Nov. 4, 2013, pp. 1-7.

Final Rejection in U.S. Appl. No. 13/633,563; mailed Apr. 22, 2015, pp. 1-12.

Final Rejection in U.S. Appl. No. 13/633,563; mailed Dec. 16, 2013, pp. 1-58.

Final Rejection issued in U.S. Appl. No. 14/172,151; mailed Jul. 20, 2015, pp. 1-18.

Final Rejection issued in U.S. Appl. No. 12/617,811; mailed Jan. 4, 2013, pp. 1-6.

Final Rejection issued in U.S. Appl. No. 12/617,811; mailed Jul. 31, 2015, pp. 1-15.

Final Rejection issued in U.S. Appl. No. 13/110,568; mailed Feb. 21, 2013, pp. 1-20.

Final Rejection issued in U.S. Appl. No. 13/310,118; mailed Aug. 2, 2012, pp. 1-20.

Final Rejection issued in U.S. Appl. No. 13/363,956; mailed May 20, 2014, pp. 1-16.

Final Rejection issued in U.S. Appl. No. 13/382,442; mailed Jul. 17, 2013, pp. 1-30.

Final Rejection issued in U.S. Appl. No. 13/382,442; mailed Jun. 6, 2014, pp. 1-29.

Final Rejection issued in U.S. Appl. No. 13/382,772; mailed Jun. 3, 2014, pp. 1-34.

Final Rejection issued in U.S. Appl. No. 13/382,772; mailed Feb. 10, 2015, pp. 1-36.

Final Rejection issued in U.S. Appl. No. 13/467,707; mailed Feb. 25, 2014, pp. 1-18.

Final Rejection issued in U.S. Appl. No. 13/467,707; mailed Jan. 7, 2015, pp. 1-8.

Final Rejection issued in U.S. Appl. No. 13/468,422; mailed Jan. 6, 2014, pp. 1-21.

Final Rejection issued in U.S. Appl. No. 13/468,422; mailed Jan. 23, 2015, pp. 1-27.

Final Rejection issued in U.S. Appl. No. 13/469,633; mailed Dec. 4, 2013, pp. 1-17.

Final Rejection issued in U.S. Appl. No. 13/469,633; mailed Jan. 23, 2015, pp. 1-12.

Final Rejection issued in U.S. Appl. No. 13/509,507; mailed Jul. 23, 2015, pp. 1-11.

Final Rejection issued in U.S. Appl. No. 13/509,542; mailed Nov. 21, 2013, pp. 1-34.

Final Rejection issued in U.S. Appl. No. 13/509,542, mailed Jan. 28, 2015, pp. 1-26.

Final Rejection issued in U.S. Appl. No. 13/595,590; mailed Apr. 2, 2014, pp. 1-7.

Final Rejection issued in U.S. Appl. No. 13/595,590; mailed Dec. 19, 2014, pp. 1-14.

Final Rejection issued in U.S. Appl. No. 13/661,476, mailed on Oct. 2, 2014, pp. 1-33.

Final Rejection issued in U.S. Appl. No. 13/692,640; mailed Jan. 6, 2015, pp. 1-11.

Final Rejection issued in U.S. Appl. No. 13/692,640; mailed Jun. 2, 2015, pp. 1-12.

Final Rejection issued in U.S. Appl. No. 13/700,631; mailed Apr. 13, 2015, pp. 1-19.

Final Rejection issued in U.S. Appl. No. 13/700,631; mailed Jun. 18, 2014, pp. 1-25.

Final Rejection issued in U.S. Appl. No. 13/819,114; mailed Mar. 2, 2015, pp. 1-10.

Final-Rejection issued in U.S. Appl. No. 13/432,811; mailed Dec. 12, 2014, pp. 1-8.

Final-Rejection issued in U.S. Appl. No. 13/432,811; mailed Mar. 31, 2015, pp. 1-9.

Final-Rejection issued in U.S. Appl. No. 13/432,811; mailed Sep. 6, 2014, pp. 1-8.

Fonseca V.A., et al., “Efficacy and Safety of the once-Daily GLP-1 Receptor Agonist Lixisenatide in Monotherapy: A Randomized, Double-Blind, Placebo-Controlled Trial in Patients with Type 2 Diabetes (Getgoal-Mono),” Diabetes Care, 2012, vol. 35 (6), pp. 1225-1231.

Fox J.D., et al., “Single Amino Acid Substitutions on the Surface of *Escherichia coli* Maltose-Binding Protein can have a Profound Impact on the Solubility of Fusion Proteins,” Protein Science, 2001, vol. 10 (3), pp. 622-630.

Fransson J., et al., “Oxidation of Human Insulin-Like Growth Factor I in formulation Studies: Kinetics of Methionine Oxidation in Aqueous Solution and in Solid State,” Pharmaceutical Research, 1996, vol. 13 (8), pp. 1252-1257.

Galloway J.A., et al., “New forms of Insulin,” Diabetes, 1972, vol. 21 (2 Suppl), pp. 637-648.

Gallwitz B., “Liraglutide. GLP-1 Receptor Agonist Treatment of Type 2 Diabetes Treatment of Obesity,” Drugs of the Future, 2008, vol. 33 (1), pp. 13-20.

Gandhi S., et al., “Molecular Pathogenesis of Parkinson’s Disease,” Human Molecular Genetics, 2005, vol. 14 (18), pp. 2749-2755.

Garber A., et al., “Liraglutide Versus Glimepiride Monotherapy for Type 2 Diabetes (Lead-3 Mono): A Randomised, 52-Week, Phase III, Double-Blind, Parallel-Treatment Trial,” The Lancet, 2009, vol. 373 (9662), pp. 473-481.

Garg R., et al., “U-500 Insulin: Why, When and How to Use in Clinical Practice,” Diabetes/Metabolism Research and Reviews, 2007, vol. 23 (4), pp. 265-268.

Garriques L.N., et al., “The Effect of Mutations on the Structure of Insulin Fibrils Studied by Fourier Transform infrared (FTIR) Spectroscopy and Electron Microscopy,” Journal of Pharmaceutical Sciences, 2002, vol. 91 (12), pp. 2473-2480.

Gault V.A., et al., “GLP-1 Agonists Facilitate Hippocampal LTP and Reverse the Impairment of LTP induced by Beta-Amyloid,” European Journal of Pharmacology, 2008, vol. 587 (1-3), pp. 112-117.

Gavin J.R., “Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus,” Diabetes Care, 1997, vol. 20 (7), pp. 1183-1197.

Geiger R., “The Chemistry of Insulin,” Chemiker Zeitung, 1976, vol. 100 (3), pp. 54-56.

Gengler S., et al., “Val(8)GLP-1 Rescues Synaptic Plasticity and Reduces Dense Core Plaques in APP/PS1 Mice,” Neurobiology of Aging, 2012, vol. 33 (2), pp. 265-276.

US 9,526,764 B2

Page 10

(56)

References Cited

OTHER PUBLICATIONS

Gerich et al., "Monotherapy with GLP-1 receptor agonist, Lixisenatide, significantly improves glycaemic control in type 2 diabetic patients," *Diabetologia* 53(Supplement 1)p. S330, Abstract 830, Presented at 46th Annual Meeting of EASD, Stockholm, Sweden, p. 1 (Sep. 2010).

Giugliano D., et al., "Treatment Regimens with Insulin Analogues and Haemoglobin A1C Target of $\leq 7\%$ in Type 2 Diabetes: A Systematic Review," *Diabetes Research and Clinical Practice*, 2010, vol. 92 (1), pp. 1-10.

Goke R., et al., "Distribution of GLP-1 Binding Sites in the Rat Brain: Evidence That Exendin-4 is a Ligand of Brain GLP-1 Binding Sites," *European Journal of Neuroscience*, 1995, vol. 7 (11), pp. 2294-2300.

Goldstein D.E., et al., "Tests of Glycemia in Diabetes," *Diabetes Care*, 1995, vol. 18 (6), pp. 896-909.

Gough K., et al., "Assessment of Dose Proportionality: Report from the Statisticians in the Pharmaceutical Industry/ Pharmacokinetics UK Joint Working Party," *Drug Information Journal*, 1995, vol. 29, pp. 1039-1048.

Goykhman S., et al., "Insulin Glargine: A Review 8 Years After its introduction," *Expert Opinion on Pharmacotherapy*, 2009, vol. 10 (4), pp. 705-718.

Greig N.H., et al., "Once Daily Injection of Exendin-4 to Diabetic Mice Achieves Long-Term Beneficial Effects on Blood Glucose Concentrations," *Diabetologia*, 1999, vol. 42 (1), pp. 45-50.

Guidance for Industry, US Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research (COER), pp. 1-11, Feb. 2014.

"Guideline for Management of Postmeal Glucose," International Diabetes Federation, pp. 1-27, Lesaffre Printers, ISBN 2-930229-48-9 (2007).

Gura T., "Systems for Identifying New Drugs Are often Faulty," *Science*, 1997, vol. 278 (5340), pp. 1041-1042.

Gutniak M., et al., "Antidiabetogenic Effect of Glucagon-Like Peptide-1 (7-36) Amide in Normal Subjects and Patients with Diabetes Mellitus," *The New England Journal of Medicine*, 1992, vol. 326 (20), pp. 1316-1322.

Gygi S.P., et al., "Quantitative Analysis of Complex Protein Mixtures Using Isotope-Coded Affinity Tags," *Nature Biotechnology*, 1999, vol. 17 (10), pp. 994-999.

Hamilton A., et al., "Novel GLP-1 Mimetics Developed to Treat Type 2 Diabetes Promote Progenitor Cell Proliferation in the Brain," *Journal of Neuroscience Research*, 2011, vol. 89 (4), pp. 481-489.

Hamilton A., et al., "Receptors for the incretin Glucagon-Like Peptide-1 are Expressed on Neurons in the Central Nervous System," *NeuroReport*, 2009, vol. 20 (13), pp. 1161-1166.

Hanas R., et al., "2010 Consensus Statement on the Worldwide Standardization of the Hemoglobin A1C Measurement," *Diabetes Care*, 2010, vol. 33 (8), pp. 1903-1904.

Hanefeld M., et al., "The Postprandial State and the Risk of Atherosclerosis," *Diabetic Medicine*, 1997, vol. 14 (Suppl 3), pp. S6-S11.

Hanefeld M., "Normnahe Postprandiale Hyperglykamie-Eine Essenzielle Komponente Guter Diabeteskontrolle Und Pravention Kardiovaskularer Erkrankungen (Near-Normal Postprandial Hyperglycemia—An Essential Component of Good Diabetes Control and Prevention of Cardiovascular Diseases)," *Paul Langerhans Lecture Diabetologie und Stoffwechsel*, 2007, vol. 2, pp. 362-369.

Hanna et al., "Canadian Diabetes Association Clinical Practice Guidelines Expert Committee Pharmacologic Management of Type 2 Diabetes," *Canadian Journal of Diabetes*, 2003, vol. 27 (Supp 2), pp. S37-S42.

Harkavyi A., et al., "Glucagon-Like Peptide I Receptor Stimulation Reverses Key Deficits in Distinct Rodent Models of Parkinson's Disease," *Journal of Neuroinflammation*, 2008, vol. 5 (19), pp. 1-9.

Harris S.B., et al., "Clinical inertia in Patients with T2Dm Requiring Insulin in Family Practice," *Canadian Family Physician*, 2010, vol. 56 (12), pp. e418-e424.

Hartmann H., et al., "Biological Activity of Des-(B26-B30)-Insulinamide and Related Analogues in Rat Hepatocyte Cultures," *Diabetologia*, 1989, vol. 32 (7), pp. 416-420.

Hellstrom M., et al., "T1388 GTP-010 as a Therapeutic tool In IBS Pain Relief: Prospective, Randomized, Placebo-Controlled Study of a GLP-1 Analog," *Gastroenterology*, 2008, vol. 134 (4), pp. A-544.

Higgins G.C., et al., "Oxidative Stress: Emerging Mitochondrial and Cellular themes and Variations in Neuronal Injury," *Journal of Alzheimer's Disease*, 2010, vol. 20, pp. S453-S473.

Himeno T., et al., "Beneficial Effects of Exendin-4 on Experimental Polyneuropathy in Diabetic Mice," *Diabetes*, 2011, vol. 60 (9), pp. 2397-2406.

Hinds K., et al., "Synthesis and Characterization of Poly(Ethylene Glycol)-Insulin Conjugates," *Bioconjugate Chemistry*, 2000, vol. 11 (2), pp. 195-201.

Hoe 901/2004 Study Investigators Group, "Safety and Efficacy of Insulin Glargine (Hoe 901) Versus NPH Insulin in Combination with oral Treatment in Type 2 Diabetic Patients," *Diabetic Medicine*, 2003, vol. 20, pp. 545-551.

Holman R.R., et al., "10-Year Follow-Up of intensive Glucose Control in Type 2 Diabetes," *The New England Journal of Medicine*, 2008, vol. 359 (15), pp. 1577-1589.

Holscher C., "Development of Beta-Amyloid-induced Neurodegeneration in Alzheimer's Disease and Novel Neuroprotective Strategies," *Reviews in the Neurosciences*, 2005, vol. 16 (3), pp. 181-212.

Holscher C., et al., "New Roles for Insulin-Like Hormones in Neuronal Signalling and Protection: New Hopes for Novel Treatments of Alzheimer's Disease?," *Neurobiology of Aging*, 2008, vol. 31 (9), pp. 1495-1502.

Holscher C., "Incretin Analogues that have been Developed to Treat Type 2 Diabetes Hold Promise as a Novel Treatment Strategy for Alzheimer's Disease," *Recent Patents on Cns Drug Discovery*, 2010, vol. 5 (2), pp. 109-117.

Holscher C., "Possible Causes of Alzheimer's Disease: Amyloid Fragments, Free Radical, and Calcium Homeostasis," *Neurobiology of Disease*, 1998, vol. 5 (3), pp. 129-141.

Holscher C., "The Role of GLP-1 in Neuronal Activity and Neurodegeneration," *Vitamins and Hormones*, 2010, vol. 84, pp. 331-354.

Holst J.J., et al., "Combining GLP-1 Receptor Agonists with Insulin: therapeutic Rationales and Clinical Findings," *Diabetes, Obesity and Metabolism*, 2013, vol. 15 (1), pp. 3-14.

Home P.D., et al., "Insulin Treatment: A Decade of Change," *British Medical Bulletin*, 1989, vol. 45 (1), pp. 92-110.

<http://diabetes.emedtv.com/lantus/generic-lantus.html>; one page, last accessed Dec. 23, 2015.

Humalog® prescribing information, Apr. 2012, pp. 1-6.

Hunter K., et al., "Drugs Developed to Treat Diabetes, Liraglutide and Lixisenatide, Cross the Blood Brain Barrier and Enhance Neurogenesis," *BMC Neuroscience*, 2012, vol. 13, pp. 1-6.

Inpharma, Product News. "AVE0010 set to deliver in type 2 diabetes mellitus," *Database Adisnews*, retrieved from STN, Jun. 2008, pp. 1-3.

"Insulin Aspart Injection." Formulated Preparations: Specific Monographs. *British Pharmacopoeia* 3. pp. 1-3 (2012).

International Search Report by the ISA for International Application No. PCT/EP2007/005932; dated Oct. 9, 2007, pp. 1-6.

International Search Report by the ISA for International Application No. PCT/EP2009/000017; dated Jun. 22, 2009, pp. 1-7.

International Search Report by the ISA for International Application No. PCT/EP2010/059436; dated Jun. 17, 2011, pp. 1-5.

International Search Report by the ISA for International Application No. PCT/EP2010/059438; dated Oct. 4, 2010, pp. 1-3.

International Search Report by the ISA for International Application No. PCT/EP2010/062638; Dated Mar. 18, 2011, pp. 1-5.

International Search Report by the ISA for International Application No. PCT/EP2010/067250; dated Mar. 23, 2011, pp. 1-3.

International Search Report by the ISA for International Application No. PCT/EP2011/058079; dated Mar. 22, 2012, pp. 1-6.

International Search Report by the ISA for International Application No. PCT/EP2011/058764; dated Jun. 30, 2011, pp. 1-9.

US 9,526,764 B2

Page 11

(56)

References Cited

OTHER PUBLICATIONS

International Search Report by the ISA for International Application No. PCT/EP2012/051670; dated Mar. 26, 2012, pp. 1-16.

International Search Report by the ISA for International Application No. PCT/EP2012/055660; dated May 10, 2012, pp. 1-6.

International Search Report by the ISA for International Application No. PCT/EP2012/058745; dated Jul. 12, 2012, pp. 1-6.

International Search Report by the ISA for International Application No. PCT/EP2012/058747; dated Jul. 8, 2012, pp. 1-6.

International Search Report by the ISA for International Application No. PCT/EP2012/058749; dated Jul. 31, 2012, pp. 1-6.

International Search Report by the ISA for International Application No. PCT/EP2012/058779; dated Aug. 28, 2012, pp. 1-5.

International Search Report by the ISA for International Application No. PCT/EP2012/066617; dated Nov. 22, 2012, pp. 1-5.

International Search Report by the ISA for International Application No. PCT/EP2012/067144; dated Aug. 11, 2012, pp. 1-5.

International Search Report by the ISA for International Application No. PCT/EP2012/069483; dated Nov. 29, 2011, pp. 1-4.

International Search Report by the ISA for International Application No. PCT/EP2012/069485; dated Dec. 11, 2012, pp. 1-5.

International Search Report by the ISA for International Application No. PCT/EP2012/071271; dated Jan. 30, 2013, pp. 1-5.

International Search Report by the ISA for International Application No. PCT/EP2012/074150; dated Nov. 20, 2012, pp. 1-4.

International Search Report by the ISA for International Application No. PCT/EP2014/051976; dated Mar. 4, 2014, pp. 1-3.

International Search Report by the ISA for International Application No. PCT/EP2014/056498; dated Jun. 25, 2014, pp. 1-10.

International Search Report by the ISA for International Application No. PCT/EP2014/062418; dated Sep. 22, 2014, pp. 1-4.

Inzucchi S.E., et al., "Management of Hyperglycemia in Type 2 Diabetes: A Patient-Centered Approach," *Diabetes Care*, 2012, vol. 35, pp. 1364-1379.

Isacson R., et al., "The Glucagon-Like Peptide 1 Receptor Agonist Exendin-4 improves Reference Memory Performance and Decreases Immobility in the forced Swim Test," *European Journal of Pharmacology*, 2009, vol. 650 (1), pp. 249-255.

ISPAD, International Diabetes Federation; "Global/IDF/ISPAD Guideline for Diabetes in Childhood and Adolescence," pp. 1-132 (2011).

Jackson R.L., et al., "Neutral Regular Insulin," *Diabetes*, 1972, vol. 21 (4), pp. 235-245.

Jain R.K., "Barriers to Drug Delivery in Solid Tumors," *Scientific American*, 1994, vol. 271 (1), pp. 58-65.

Jang J.H., et al., "Neuroprotective Effects of Triticum Aestivum L. Against B-Amyloid-induced Cell Death and Memory Impairments," *Phytotherapy Research*, 2010, vol. 24 (1), pp. 76-84.

Jekel P.A., et al., "Use of Endoproteinase Lys-C from *Lysobacter* Enzymogenes in Protein Sequence Analysis," *Analytical Biochemistry*, 1983, vol. 134 (2), pp. 347-354.

Jendle J., et al., "Insulin and GLP-1 Analog Combinations in Type 2 Diabetes Mellitus: A Critical Review," *Expert Opinion on Investigational Drugs*, 2012, vol. 21 (10), pp. 1463-1474.

Jimenez S., et al., "Inflammatory Response in the Hippocampus of PS1M146L/App751SL Mouse Model of Alzheimer's Disease: Age-Dependent Switch in the Microglial Phenotype from Alternative to Classic," *The Journal of Neuroscience*, 2008, vol. 28 (45), pp. 11650-11661.

Johnson et al., "When is a unit of insulin not a unit of insulin? Detemir dosing in type 2 diabetes" Poster, one page, 2008. <http://professional.diabetes.org/ContentUPosters/2008/p8-LB.pdf>.

Jorgensen K.H., et al., "Five Fold Increase of Insulin Concentration Delays the Absorption of Subcutaneously Injected Human Insulin Suspensions in Pigs," *Diabetes Research and Clinical Practice*, 2000, vol. 50, pp. 161-167.

Kaarsholm N.C., et al., "Engineering Stability of the Insulin Monomer Fold with Application to Structure-Activity Relationships," *Biochemistry*, 1993, vol. 32 (40), pp. 10773-10778.

Kadima W., "Role of Metal Ions in the T- to R-Allosteric Transition in the Insulin Hexamer," *Biochemistry*, 1999, vol. 38 (41), pp. 13443-13452.

Kaduszkiewicz H., et al., "Cholinesterase inhibitors for Patients with Alzheimer's Disease: Systematic Review of Randomised Clinical Trials," *British Medical Journal (Clinical Research ed.)*, 2005, vol. 331 (7512), pp. 321-327.

Kaech S., et al., "Culturing Hippocampal Neurons," *Nature Protocols*, 2006, vol. 1 (5), pp. 2406-2415.

Kahn S.E., et al., "Glycemic Durability of Rosiglitazone, Metformin, or Glyburide Monotherapy," *The New England Journal of Medicine*, 2006, vol. 355 (23), pp. 2427-2443.

Kakhki V.R.D., et al., "Normal Values of Gallbladder Ejection Fraction Using 99m Tc-Sestamibi Scintigraphy after a Fatty Meal formula," *Journal of Gastrointestinal and Liver Diseases*, 2007, vol. 16 (2), pp. 157-161.

Kamisawa T., et al., "Pancreatographic investigation of Pancreatic Duct System and Pancreaticobiliary Malformation," *Journal of Anatomy*, 2008, vol. 212 (2), pp. 125-134.

Kanazawa M., et al., "Criteria and Classification of Obesity in Japan and Asia-Oceania," *Asia Pacific Journal of Clinical Nutrition*, 2002, vol. 11 (Suppl 7), pp. S732-S737.

Kang S., et al., "Subcutaneous Insulin Absorption Explained by Insulin's Physicochemical Properties Evidence from Absorption Studies of Soluble Human Insulin and Insulin Analogues in Humans," *Diabetes Care*, 1991, vol. 14 (11), pp. 942-948.

Kao C.H., et al., "The Evaluation of Gallbladder Function by Quantitative Radionuclide Cholescintigraphy in Patients with Noninsulin-Dependent Diabetes Mellitus," *Nuclear Medicine Communications*, 1993, vol. 14 (10), pp. 868-872.

Kapitza et al., Abstract "Pharmacodynamic Characteristics of Lixisenatide QD vs Liraglutide QD in Patients with T2DM Inadequately Controlled with Metformin" Abstract D-0740, presented at the World Diabetes Congress in Dubai, Dec. 5-8, 2011, one page.

Kastin A.J., et al., "Entry of Exendin-4 into Brain Is Rapid but may be Limited at High Doses International Journal of Obesity and Related Metabolic Disorders," *Journal of the International Association for the Study of Obesity*, 2003, vol. 27 (3), pp. 313-318.

Kastin A.J., et al., "Interactions of Glucagon-Like Peptide-1 (GLP-1) with the Blood-Brain Barrier," *Journal of Molecular Neuroscience*, 2001, vol. 18 (1-2), pp. 7-14.

Kemmler W., et al., "Studies on the Conversion of Proinsulin to Insulin," *The Journal of Biological Chemistry*, 1971, vol. 246 (22), pp. 6786-6791.

Kendall D.M., et al., "Effects of Exenatide (Exendin-4) on Glycemic Control Over 30 Weeks in Patients with Type 2 Diabetes Treated with Metformin and a Sulfonyleurea," *Diabetes care*, 2005, vol. 28 (5), pp. 1083-1091.

Kielgast U., et al., "Treatment of Type 1 Diabetic Patients with Glucagon-Like Peptide-1 (GLP-1) and GLP-1R Agonists," *Current Diabetes Reviews*, 2009, vol. 5 (4), pp. 266-275.

Kim S., et al., "Exendin-4 Protects Dopaminergic Neurons by Inhibition of Microglial Activation and Matrix Metalloproteinase-3 Expression in an Animal Model of Parkinson's Disease," *Journal of Endocrinology*, 2009, vol. 202 (3), pp. 431-439.

Knee T.S., et al., "A Novel Use of U-500 Insulin for Continuous Subcutaneous Insulin infusion in Patients with Insulin Resistance: A Case Series," *Endocrine Practice*, 2003, vol. 9 (3), pp. 181-186.

Knudsen L.B., et al., "Potent Derivatives of Glucagon-Like Peptide-1 with Pharmacokinetic Properties Suitable for once Daily Administration," *Journal of Medicinal Chemistry*, 2000, vol. 43 (9), pp. 1664-1669.

Kohn W.D., et al., "Pi-Shifted Insulin Analogs with Extended in Vivo Time Action and Favorable Receptor Selectivity," *Peptides*, 2007, vol. 28 (4), pp. 935-948.

Kohner E.M., "Diabetic Retinopathy," *British Medical Bulletin*, 1989, vol. 45 (1), pp. 148-173.

Kolterman O.G., et al., "Synthetic Exendin-4 (Exenatide) Significantly Reduces Postprandial and Fasting Plasma Glucose in Subjects with Type 2 Diabetes," *The Journal of Clinical Endocrinology & Metabolism*, 2003, vol. 88 (7), pp. 3082-3089.

US 9,526,764 B2

Page 12

(56)

References Cited

OTHER PUBLICATIONS

Korczyn A.D., et al, “Emerging therapies in the Pharmacological Treatment of Parkinson’s Disease,” *Drugs*, 2002, vol. 62 (5), pp. 775-786.

Krishnamurthy G.T., et al., “Constancy and Variability of Gallbladder Ejection Fraction: Impact on Diagnosis and therapy,” *Journal of Nuclear Medicine*, 2004, vol. 45 (11), pp. 1872-1877.

Lando, “The New ‘Designer’ Insulins”, *Clinical Diabetes*, 18(4): Fall 2000 (http://journal.diabetes.org/clinical_diabetesN18N42000/pg154.htm; accessed Oct. 22, 2013, pp. 1-13).

Langston J.W., et al., “Chronic Parkinsonism in Humans due to a Product of Meperedine-Analog Synthesis,” *Science*, 1983, vol. 219 (4587), pp. 979-980.

Langui D., et al., “Subcellular Topography of Neuronal A β Peptide in APPxPS1 Transgenic Mice,” *The American Journal of Pathology*, 2004, vol. 165 (5), pp. 1465-1477.

Lantus® ANNEX I—Summary of product characteristics. Date of first authorisation: Jun. 9, 2000, pp. 1-164.

Lantus® prescribing information, May 2012, pp. 1-6.

Lantus® Product Information—European Medicines Agency, first published Aug. 5, 2009, pp. 1-2.

Larsen B.D., et al., “Sequence-Assisted Peptide Synthesis (SAPS),” *Journal of Peptide Research*, 1998, vol. 52 (6), pp. 470-476.

Larsen P.J., et al., “Combination of the Insulin Sensitizer, Pioglitazone, and the Long-Acting Glp-1 Human Analog, Liraglutide, Exerts Potent Synergistic Glucose-Lowering Efficacy in Severely Diabetic ZDF Rats,” *Diabetes, Obesity and Metabolism*, 2008, vol. 10, pp. 301-311.

Laursen K., et al., “Enhanced Monitoring of Biopharmaceutical Product Purity Using Liquid Chromatography-Mass Spectrometry,” *Journal of Chromatography A*, 2011, vol. 1218 (28), pp. 4340-4348.

Lee C.H., et al., “Ischemia-Induced Changes in Glucagon-Like Peptide-1 Receptor and Neuroprotective Effect of its Agonist, Exendin-4, in Experimental Transient Cerebral Ischemia,” *Journal of Neuroscience Research*, 2011, vol. 89 (7), pp. 1103-1113.

Leib R.D., et al., “Direct Quantitation of Peptide Mixtures without Standards Using Clusters formed by Electrospray Ionization Mass Spectrometry,” *Analytical Chemistry*, 2009, vol. 81 (10), pp. 3965-3972.

Lens J., “The Terminal Carboxyl Groups of Insulin,” *Biochimica et Biophysica Acta*, 1949, vol. 3, pp. 367-370.

Levemir® prescribing information, Dec. 2011, pp. 1-6.

Levene P.A., et al., “Calculation of Isoelectric Point,” *The Journal of Biological Chemistry*, 1923, vol. 55, pp. 801-813.

Levin P., et al., “Combination therapy with Insulin Glargine and Exenatide: Real-World Outcomes in Patients with Type 2 Diabetes,” *Current Medical Research and Opinion*, 2012, vol. 28 (3), pp. 439-446.

Levine R.L., et al., “Oxidation of Methionine in Proteins: Roles in Antioxidant Defense and Cellular Regulation,” *IUBMB life*, 2000, vol. 50 (4-5), pp. 301-307.

Leyer S., et al., “The Role of the C-Terminus of the Insulin B-Chain in Modulating Structural and Functional Properties of the Hormone,” *International Journal of Peptide and Protein Research*, 1995, vol. 46 (5), pp. 397-407.

Li H., et al., “Chronic Treatment of Exendin-4 Affects Cell Proliferation and Neuroblast Differentiation in the Adult Mouse Hippocampal Dentate Gyrus,” *Neuroscience letters*, 2010, vol. 19, pp. 1205-1219.

Li L., et al., “Common Pathological Processes in Alzheimer Disease and Type 2 Diabetes: A Review,” *Brain Research Reviews*, 2007, vol. 56, pp. 384-402.

Li Y., et al., “Enhancing the GLP-1 Receptor Signaling Pathway Leads to Proliferation and Neuroprotection in Human Neuroblastoma Cells,” *Journal of Neurochemistry*, 2010, vol. 113 (6), pp. 1621-1631.

Li Y., et al., “GLP-1 Receptor Stimulation Preserves Primary Cortical and Dopaminergic Neurons in Cellular and Rodent Models

of Stroke and Parkinsonism,” *Proceedings of the National Academy of Sciences of the United States of America*, 2009, vol. 106 (4), pp. 1285-1290.

Li Y., et al., “GLP-1 Receptor Stimulation Reduces Amyloid-Beta Peptide Accumulation and Cytotoxicity in Cellular and Animal Models of Alzheimer’s Disease,” *Journal of Alzheimer’s Disease*, 2010, vol. 19 (4), pp. 1205-1219.

Lill N., “Production of Fast-Acting Insulins and Delayed-Release Insulins- How can this Problem be Solved by Technology? Insulin formulations,” *Pharmazie in Unserer Zeit*, 2001, vol. 30 (1), pp. 56-61.

Liu & Ruus, Abstract “Pharmacokinetics and Safety of the GLP-1 Agonist AVE0010 in Patients with Renal Impairment,” *Diabetes* 58 (Suppl. 1): Abstract 557-P for the 69th Scientific Session of the American Diabetes Association Jun. 5-9, 2009, New Orleans, Louisiana, pp. 1-2.

Lixisenatide, Chemical Structure CID 16139342, Pubchem, accessed Feb. 5, 2015 at URL pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?sid=135267128&viewopt=Deposited, pp. 1-3.

Insulinpraparat Wikipedia, <http://de.wikipedia.org/wiki/Insulinpr%C3%A4parat>, pp. 1-15 (Feb. 5, 2013).

Insuman® Comb25 prescribing information, Feb. 2011, pp. 1-4.

Insuman® Infusat prescribing information, Feb. 2011, pp. 1-4.

Lotharius J., et al., “Effect of Mutant Alpha-Synuclein on Dopamine Homeostasis in a New Human Mesencephalic Cell Line,” *The Journal of Biological Chemistry*, 2002, vol. 277 (41), pp. 38884-38894.

Lotharius J., et al., “Progressive Degeneration of Human Mesencephalic Neuron-Derived Cells Triggered by Dopamine-Dependent Oxidative Stress is Dependent on the Mixed-Lineage Kinase Pathway,” *Journal of Neuroscience*, 2005, vol. 25 (27), pp. 6329-6342.

Lougheed W.D., et al., “Physical Stability of Insulin Formulations,” *Diabetes*, 1983, vol. 32 (5), pp. 424-432.

Lyxumia 10 micrograms solution for injection, Summary of Product Characteristics, updated Oct. 31, 2014, pp. 1-12.

Lyxumia® Annex I—Summary of product characteristics. Date of first authorisation: Feb. 1, 2013, pp. 1-92.

Lyxumia, Chemical Subgroup A10BX, Community Register of Medicinal Products for Human Use, European Commission Public Health, p. 1-2 (May 2, 2013).

Lyxumia® Product Information—European Medicines Agency, first published Mar. 14, 2013, pp. 1-2.

Mancuso M., et al., “Clinical Features and Pathogenesis of Alzheimer’s Disease: involvement of Mitochondria and Mitochondrial DNA,” *Advances in Experimental Medicine and Biology*, 2010, vol. 685, pp. 34-44.

Marbury T.C., et al., “A Pilot Study to Examine the Feasibility of Insulin Glargine in Subjects with Impaired Fasting Glucose, Impaired Glucose tolerance or New-onset Type 2 Diabetes,” *Experimental and Clinical Endocrinology & Diabetes*, 2008, vol. 116 (5), pp. 282-288.

Margolis R.L., et al., “Diagnosis of Huntington Disease,” *Clinical Chemistry*, 2003, vol. 49 (10), pp. 1726-1732.

Markussen J., et al., “Soluble, Prolonged-Acting Insulin Derivatives. I. Degree of Protraction and Crystallizability of Insulins Substituted in the Termini of the B-Chain,” *Protein Engineering*, 1987, vol. 1 (3), pp. 205-213.

Markussen J., et al., “Soluble, Prolonged-Acting Insulin Derivatives II Degree of Protraction and Crystallizability of Insulins Substituted in Positions A17, B8, B13, B27 and B30,” *Protein Engineering*, 1987, vol. 1 (3), pp. 215-223.

Markussen J., et al., “Soluble, Prolonged-Acting Insulin Derivatives. III. Degree of Protraction, Crystallizability and Chemical Stability of Insulins Substituted in Positions A21, B13, B23, B27 and B30,” *Protein Engineering*, 1988, vol. 2 (2), pp. 157-166.

Martin B., et al., “Exendin-4 Improves Glycemic Control, Ameliorates Brain and Pancreatic Pathologies, and Extends Survival in a Mouse Model of Huntington’s Disease,” *Diabetes*, 2009, vol. 58 (2), pp. 318-328.

US 9,526,764 B2

Page 13

(56)

References Cited**OTHER PUBLICATIONS**

Martin L.J., et al., "Neurodegeneration in Excitotoxicity, Global Cerebral Ischemia, and Target Deprivation: A Perspective on the Contributions of Apoptosis and Necrosis," *Brain Research Bulletin*, 1998, vol. 46 (4), pp. 281-309.

Mattson M.P., "Calcium and Neurodegeneration," *Aging Cell*, 2007, vol. 6 (3), pp. 337-350.

McClellan P.L., et al., "Glucagon-Like Peptide-1 Analogues Enhance Synaptic Plasticity in the Brain: A Link between Diabetes and Alzheimer's Disease," *European Journal of Pharmacology*, 2010, vol. 630 (1-3), pp. 158-162.

McClellan P.L., et al., "The Diabetes Drug Liraglutide Prevents Degenerative Processes in a Mouse Model of Alzheimer's Disease," *The Journal of Neuroscience*, 2011, vol. 31 (17), pp. 6587-6594.

Mecklenburg R.S., et al., "Complications of Insulin Pump therapy: The Effect of Insulin Preparation," *Diabetes Care*, 1985, vol. 8 (4), pp. 367-370.

Medline Plus, "Obesity" available at <http://www.nlm.nih.gov/medlineplus/obesity.html>, Retrieved Aug. 22, 2013, one page.

Meier J.J., "GLP-1 Receptor Agonists for individualized Treatment of Type 2 Diabetes Mellitus," *Nature Reviews. Endocrinology*, 2012, vol. 8 (12), pp. 728-742.

Merrifield B., "Solid Phase Synthesis," *Science*, 1986, vol. 232 (4748), pp. 341-347.

Mikhail N.E., "Is Liraglutide a Useful Addition to Diabetes therapy?," *Endocrine Practice*, 2010, vol. 16 (6), pp. 1028-1037.

Monnier L., et al., "The Loss of Postprandial Glycemic Control Precedes Stepwise Deterioration of Fasting with Worsening Diabetes," *Diabetes Care*, 2007, vol. 30 (2), pp. 263-269.

Moreno-Gonzalez I., et al., "Extracellular Amyloid-B and Cytotoxic Glial Activation Induce Significant Entorhinal Neuron Loss in Young PS1M146L/APP751SL Mice," *Journal of Alzheimer's Disease*, 2009, vol. 18, pp. 755-776.

Moretto T.J., et al., "Efficacy and tolerability of Exenatide Monotherapy Over 24 Weeks in Antidiabetic Drug-Naive Patients with Type 2 Diabetes: A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study," *Clinical Therapeutics*, 2008, vol. 30 (8), pp. 1448-1460.

Muller G., et al., "Insulin Signaling in the Yeast *Saccharomyces Cerevisiae*. 1. Stimulation of Glucose Metabolism and Snf 1 Kinase by Human Insulin," *Biochemistry*, 1998, vol. 37 (24), pp. 8683-8695.

Muzaffar M., et al., "The Mechanism of Enhanced Insulin Amyloid Fibril formation by Naciis Better Explained by a Conformational Change Model," *PLoS One*, 2011, vol. 6 (11), pp. 1-11.

Nakagawa A., et al., "Receptor Gene Expression of Glucagon-Like Peptide-1, but not Glucose-Dependent Insulinotropic Polypeptide, in Rat Nodose Ganglion Cells," *Autonomic Neuroscience*, 2004, vol. 110, pp. 36-43.

Nathan D.M., et al., "Management of Hyperglycaemia in Type 2 Diabetes Mellitus: A Consensus Algorithm for the initiation and Adjustment of therapy. Update Regarding the Thiazolidinediones," *Diabetologia*, 2008, vol. 51 (1), pp. 8-11.

Nathan M.D., et al., "Insulinotropic Action of Glucagon Like Peptide-1-(7-37) in Diabetic and Nondiabetic Subjects," *Diabetes Care*, 1992, vol. 15 (2), pp. 270-276.

Nauck M.A., et al., "Comparative Evaluation of Incretin-Based Antidiabetic Medications and Alternative therapies to be Added to Melformin in the Case of Monotherapy Failure," *Journal of Diabetes Investigation*, 2010, vol. 1 (1-2), pp. 24-36.

Nauck M.A., et al., "Effects of Subcutaneous Glucagon-Like Peptide 1 (GLP-1 [7-36 Amide]) in Patients with NIDDM," *Diabetologia*, 1996, vol. 39 (12), pp. 1546-1553.

Nauck M.A., et al., "Glucagon-Like Peptide 1 and its Potential in the Treatment of Non-Insulin-Dependent Diabetes Mellitus," *Hormone and Metabolic Research*, 1997, vol. 29 (9), pp. 411-416.

NCT00299871, ClinicalTrials.gov, "Dose Ranging Study of the GLP-1 Agonist AVE0010 in Metformin-Treated Subjects With Type 2 Diabetes Mellitus," Jun. 22, 2010, Retrieved Nov. 7, 2011, pp. 1-5.

NCT00688701 Clinical Trials.gov "GLP-1 Receptor Agonist Lixisenatide in Patients with Type 2 Diabetes for Glycemic Control and Safety Evaluation in Monotherapy (GETGOAL-MONO)" accessed Jul. 27, 2014; pp. 1-5.

NCT00712673, Clinical Trials.gov, "GLP-A Agonist AVE0010 (Morning or Evening) in Patients with Type 2 Diabetes for Glycemic Control and Safety Evaluation, on Top of Metformin", Mar. 22, 2011, pp. 1-4.

NCT00715624 Clinical Trials.gov "GLP-1 Agonist AVE0010 in Patients With Type 2 Diabetes for Glycemic Control and Safety Evaluation, on Top of Basal Insulin" (updated Jul. 20, 2008), p. 1-3.

NCT00715624 Clinical Trials.gov "GLP-1 Receptor Agonist Lixisenatide in Patients With Type 2 Diabetes for Glycemic Control and Safety Evaluation, on Top of Basal Insulin (GETGOAL-L)" (2008-2014), p. 1-6 (Feb. 2011).

Final Rejection issued in U.S. Appl. No. 13/382,442; mailed Aug. 11, 2015, pp. 1-35.

NCT00763815, ClinicalTrials.gov, U.S. National Institutes of Health: "GLP-1 Agonist AVE0010 in Patients With Type 2 Diabetes for Glycemic Control and Safety Evaluation on Top of Pioglitazone (GETGOAL-P)" pp. 1-8 (Jun. 27, 2011).

NCT00975286, Clinical Trials.gov, "24-week Treatment with Lixisenalide in Type 2 Diabetes Insufficiently Controlled With Melformin and Insulin Glargine", Aug. 8, 2011, pp. 1-4.

NCT00976937, ClinicalTrials.gov, "24-week Study Comparing Lixisenatide (AVE0010) to Sitagliptin as add-on to Metformin in Obese Type 2 Diabetic Patients Younger Than 50," May 6, 2011, Retrieved Nov. 7, 2011, pp. 1-4.

NCT01146678, ClinicalTrials.gov "Relative Bioavailability and Activity of Different Formulations of Insulin Glargine and Lixisenatide in Patients With Diabetes Mellitus Type 1" last updated Sep. 10, 2010, pp. 1-4.

NCT01174810, ClinicalTrials.gov "Exendin-4 as a Treatment for Parkinson's Disease—Pilot Study" accessed Aug. 8, 2011, pp. 1-5.

NCT01195454, NIH Clinical Trials, "Euglycemic clamp dose-response study comparing insulin glargine U300 with Lantus U100" last updated Dec. 13, 2010, pp. 1-4.

NCT01255163, ClinicalTrials.gov "A Clinical Trial of Exendin-4 for the Treatment of Alzheimer's Disease" accessed Aug. 8, 2011, pp. 1-7.

NCT02058147 ClinicalTrials.gov "Efficacy and Safety of Insulin Glargine/Lixisenatide Fixed Ratio Combination Compared to Insulin Glargine Alone and Lixisenatide Alone on Top Metformin in Patients With T2DM (LixLan-O)" first received by ClinicalTrials.gov on Feb. 6, 2014, pp. 1-3.

NCT02058160 ClinicalTrials.gov "Efficacy and Safety of the Insulin Glargine/Lixisenatide Fixed Ratio Combination Versus Insulin Glargine in Patients With Type 2 Diabetes (LixiLan-L)" first received by ClinicalTrials.gov on Feb. 6, 2014, pp. 1-3.

Needleman S.B., et al., "A General Method Applicable to the Search for Similarities in the Amino Acid Sequence of Two Proteins," *Journal of Molecular Biology*, 1970, vol. 48 (3), pp. 443-453.

Neidle, "18.2 Failure Modes in the Discovery Process" *Cancer Drug Design and Discovery*, Elsevier/Academic Press, pp. 427-431 (2008).

Nettleton E.J., et al., "Characterization of the Oligomeric States of Insulin in Self-Assembly and Amyloid Fibril formation by Mass Spectrometry," *Biophysical Journal*, 2000, vol. 79 (2), pp. 1053-1065.

Nicklas et al., "Inhibition of NADH-Linked Oxidation in Brain Mitochondria by 1-Methyl-4-Phenyl-Pyridine, A Metabolite of the Neurotoxin, 1-Methyl-4-Phenyl-1,2,5,6-Tetrahydropyridine," *Life Sciences*, 1985, vol. 36, pp. 2503-2508.

Nielsen L.L., et al., "Pharmacology of Exenatide (Synthetic Exendin-4): A Potential therapeutic for Improved Glycemic Control of Type 2 Diabetes," *Regulatory Peptides*, 2004, vol. 117 (2), pp. 77-88.

Noble S.L., et al., "Insulin Lispro: A Fast-Acting Insulin Analog," *American Family Physician*, 1998, vol. 57 (2), pp. 279-286.

Non-Final Office Action from U.S. Appl. No. 12/617,805; mailed Jul. 24, 2014, pp. 1-12.

Non-Final Office Action from U.S. Appl. No. 12/617,805; mailed May 2, 2012, pp. 1-11.

US 9,526,764 B2

Page 14

(56)

References Cited**OTHER PUBLICATIONS**

Non-Final Office Action from U.S. Appl. No. 12/617,805; mailed May 10, 2011, pp. 1-12.

Non-Final Office Action from U.S. Appl. No. 12/617,805; mailed Sep. 15, 2015, pp. 1-12.

Non-Final Office Action issued in U.S. Appl. No. 13/602,913; mailed on Dec. 2, 2014, pp. 1-12.

Non-Final Office Action issued in U.S. Appl. No. 13/602,913; mailed on Jan. 13, 2014, pp. 1-53.

Non-Final Office Action issued in U.S. Appl. No. 13/602,913; mailed on May 17, 2013, pp. 1-7.

Non-Final Rejection in U.S. Appl. No. 13/633,496; mailed Apr. 29, 2013, pp. 1-53.

Non-Final Rejection in U.S. Appl. No. 13/633,496; mailed Apr. 6, 2015, pp. 1-14.

Non-Final Rejection in U.S. Appl. No. 13/633,496; mailed May 22, 2014, pp. 1-11.

Non-Final Rejection in U.S. Appl. No. 13/633,563; mailed Dec. 1, 2014, pp. 1-9.

Non-Final Rejection in U.S. Appl. No. 13/633,563; mailed Jul. 1, 2013, pp. 1-56.

Non-Final Rejection in U.S. Appl. No. 13/633,563; mailed Jun. 4, 2014, pp. 1-11.

Non-Final Rejection in U.S. Appl. No. 13/633,563; mailed Oct. 6, 2015, pp. 1-12.

Non-Final Rejection issued in U.S. Appl. No. 14/172,151; mailed Mar. 24, 2015, pp. 1-16.

Non-Final Rejection issued in U.S. Appl. No. 12/617,811; mailed Apr. 27, 2011, pp. 1-10.

Non-Final Rejection issued in U.S. Appl. No. 12/617,811; mailed Jan. 14, 2015, pp. 1-15.

Non-Final Rejection issued in U.S. Appl. No. 12/617,811; mailed Jun. 21, 2012, pp. 1-7.

Non-Final Rejection issued in U.S. Appl. No. 12/617,811; mailed Oct. 27, 2011, pp. 1-13.

Non-Final Rejection issued in U.S. Appl. No. 13/110,568; mailed Mar. 19, 2012, pp. 1-18.

Non-Final Rejection issued in U.S. Appl. No. 13/310,118; mailed Mar. 19, 2012, pp. 1-19.

Non-Final Rejection issued in U.S. Appl. No. 13/310,118, mailed Mar. 29, 2013, pp. 1-21.

Non-Final Rejection issued in U.S. Appl. No. 13/310,118, mailed Mar. 25, 2015, pp. 1-15.

Non-Final Rejection issued in U.S. Appl. No. 13/363,956; mailed Apr. 10, 2013, pp. 1-15.

Non-Final Rejection issued in U.S. Appl. No. 13/363,956; mailed Feb. 11, 2015, pp. 1-18.

Non-Final Rejection issued in U.S. Appl. No. 13/363,956; mailed May 29, 2015, pp. 1-17.

Non-Final Rejection issued in U.S. Appl. No. 13/363,956; mailed Nov. 20, 2013, pp. 1-10.

Non-Final Rejection issued in U.S. Appl. No. 13/382,442; mailed Nov. 7, 2012, pp. 1-26.

Non-Final Rejection issued in U.S. Appl. No. 13/382,442; mailed Dec. 19, 2013, pp. 1-29.

Non-Final Rejection issued in U.S. Appl. No. 13/382,442; mailed Feb. 5, 2015, pp. 1-31.

Non-Final Rejection issued in U.S. Appl. No. 13/382,772; mailed Nov. 21, 2013, pp. 1-42.

Non-Final Rejection issued in U.S. Appl. No. 13/382,772; mailed Sep. 29, 2014, pp. 1-33.

Non-Final Rejection issued in U.S. Appl. No. 13/382,772; mailed Sep. 14, 2015, pp. 1-42.

Non-Final Rejection issued in U.S. Appl. No. 13/467,707; mailed Jul. 15, 2013, pp. 1-20.

Non-Final Rejection issued in U.S. Appl. No. 13/467,707; mailed Jul. 25, 2014, pp. 1-22.

Non-Final Rejection issued in U.S. Appl. No. 13/467,707; mailed Sep. 16, 2015, pp. 1-13.

Non-Final Rejection issued in U.S. Appl. No. 13/468,422; mailed Nov. 4, 2015; pp. 1-27.

Non-Final Rejection issued in U.S. Appl. No. 13/469,633; mailed Aug. 19, 2013, pp. 1-25.

Non-Final Rejection issued in U.S. Appl. No. 13/469,633; mailed Aug. 22, 2014, pp. 1-23.

Non-Final Rejection issued in U.S. Appl. No. 13/469,633; mailed Mar. 27, 2013, pp. 1-39.

Non-Final Rejection issued in U.S. Appl. No. 13/509,507; mailed Aug. 6, 2013, pp. 1-11.

Non-Final Rejection issued in U.S. Appl. No. 13/509,507; mailed Sep. 19, 2014, pp. 1-9.

Non-Final Rejection issued in U.S. Appl. No. 13/509,507; mailed Feb. 19, 2015, pp. 1-10.

Non-Final Rejection issued in U.S. Appl. No. 13/509,507; mailed Dec. 8, 2015, pp. 1-14.

Non-Final Rejection issued in U.S. Appl. No. 13/509,542; mailed May 23, 2013, pp. 1-21.

Non-Final Rejection issued in U.S. Appl. No. 13/509,542; mailed Apr. 2, 2014, pp. 1-20.

Non-Final Rejection issued in U.S. Appl. No. 13/509,542; mailed Aug. 11, 2015, pp. 1-30.

Non-Final Rejection issued in U.S. Appl. No. 13/595,590; mailed Jun. 5, 2015, pp. 1-9.

Non-Final Rejection issued in U.S. Appl. No. 13/595,590; mailed Oct. 16, 2013, pp. 1-7.

Non-Final Rejection issued in U.S. Appl. No. 13/595,590; mailed Sep. 5, 2014, pp. 1-10.

Non-Final Rejection issued in U.S. Appl. No. 13/661,476; mailed Jun. 4, 2015, pp. 1-31.

Non-Final Rejection issued in U.S. Appl. No. 13/661,476, mailed on Dec. 4, 2013, pp. 1-11.

Non-Final Rejection issued in U.S. Appl. No. 13/661,476, mailed on Mar. 6, 2014, pp. 1-28.

Non-Final Rejection issued in U.S. Appl. No. 13/661,476, mailed on Sep. 16, 2013, pp. 1-19.

Non-Final Rejection issued in U.S. Appl. No. 13/692,640; mailed May 6, 2014, pp. 1-10.

Non-Final Rejection issued in U.S. Appl. No. 13/692,640; mailed Oct. 31, 2013, pp. 1-10.

Non-Final Rejection issued in U.S. Appl. No. 13/700,631; mailed Apr. 22, 2013, pp. 1-27.

Non-Final Rejection issued in U.S. Appl. No. 13/700,631; mailed Dec. 17, 2015, pp. 1-18.

Non-Final Rejection issued in U.S. Appl. No. 13/700,631; mailed Nov. 18, 2014, pp. 1-22.

Non-Final Rejection issued in U.S. Appl. No. 13/700,631; mailed Nov. 29, 2013, pp. 1-23.

Non-Final Rejection issued in U.S. Appl. No. 13/819,114; mailed Dec. 2, 2015, pp. 1-14.

Non-Final Rejection issued in U.S. Appl. No. 13/819,114; mailed Jul. 31, 2014, pp. 1-9.

Non-Final Rejection issued in U.S. Appl. No. 14/220,562; mailed Apr. 8, 2015, pp. 1-18.

Non-Final Rejection issued in U.S. Appl. No. 14/303,895; mailed May 21, 2015, pp. 1-11.

Non-Final Rejection issued in U.S. Appl. No. 14/303,895; mailed Sep. 9, 2015, pp. 1-12.

Non-Final Rejection issued in U.S. Appl. No. 13/467,757; mailed Apr. 17, 2013, pp. 1-9.

Non-Final Rejection issued in U.S. Appl. No. 13/468,422; mailed Jun. 4, 2014, pp. 1-24.

Non-Final Rejection issued in U.S. Appl. No. 13/468,422; mailed Mar. 27, 2013, pp. 1-27.

Non-Final Rejection issued in U.S. Appl. No. 13/468,422; mailed Sep. 16, 2013, pp. 1-19.

Non-Final Rejection issued in U.S. Appl. No. 13/382,772; mailed Apr. 10, 2013, pp. 1-48.

Non-Final-Rejection issued in U.S. Appl. No. 13/432,811; mailed Apr. 8, 2013, pp. 1-7.

Non-Final-Rejection issued in U.S. Appl. No. 13/432,811; mailed Jul. 29, 2014, pp. 1-8.

US 9,526,764 B2

Page 15

(56)

References Cited

OTHER PUBLICATIONS

Non-Final-Rejection issued in U.S. Appl. No. 13/432,811; mailed Sep. 9, 2015, pp. 1-11.

Novolog® product information, Oct. 2009, pp. 1-4.

NovoMix® prescribing information, Feb. 2011, pp. 1-5.

NovoRapid® prescribing information, Jul. 2012, pp. 1-5.

Organization for Economic Co-Operation and Development; OECD Principles of Good Laboratory Practice and Compliance Monitoring (as revised in 1997); ENV/MC/CHEM (98)17:1-41 (Jan. 21, 1998).

Orskov C., "Glucagon-Like Peptide-1, A New Hormone of the Entero-insular Axis," *Diabetologia*, 1992, vol. 35 (8), pp. 701-711.

Ott P., et al., "Diabetes in Germany(Dig) Study A Prospective 4-Year-Follow-Up Study on the Quality of Treatment for Type 2 Diabetes in Daily Practice," *Deutsche Medizinische Wochenschrift*, 2009, vol. 134 (7), pp. 291-297.

Park C.W., et al., "PPARalpha Agonist Fenofibrate Improves Diabetic Nephropathy in Db/Db Mice," *Kidney International*, 2006, vol. 69 (9), pp. 1511-1517.

Parkin "Guideline for Management of Postmeal Glucose" International Diabetes Federation, pp. 1-32 (Oct. 2007).

Patel & Advance Collaborative Group, "Effects of a Fixed Combination of Perindopril and indapamide on Macrovascular and Microvascular Outcomes in Patients with Type 2 Diabetes Mellitus (the Advance Trial): A Randomised Controlled Trial," *Lancet*, 2007, vol. 370 (9590), pp. 829-840.

Patel K., et al., "Chemical Pathways of Peptide Degradation. II. Kinetics of Deamidation of an Asparaginyl Residue in a Model Hexapeptide," *Pharmaceutical Research*, 1990, vol. 7 (8), pp. 703-711.

Pederson R.A., et al., "Improved Glucose tolerance in Zucker Fatty Rats by oral Administration of the Dipeptidyl Peptidase IV inhibitor Isoleucine Thiazolidide," *Diabetes*, 1998, vol. 47 (8), pp. 1253-1258.

Perfetti R., "Combining Basal Insulin Analogs with Glucagon-Like Peptide-1 Mimetics," *Diabetes Technology & Therapeutics*, 2011, vol. 13 (9), pp. 873-881.

Perry T., et al., "A Novel Neurotrophic Property of Glucagon-Like Peptide 1: A Promoter of Nerve Growth Factor-Mediated Differentiation in PC12 Cells," *The Journal of Pharmacology and Experimental Therapeutics*, 2002, vol. 300 (3), pp. 958-966.

Perry T., et al., "Evidence of GLP-1-Mediated Neuroprotection in an Animal Model of Pyridoxine-induced Peripheral Sensory Neuropathy," *Experimental Neurology*, 2007, vol. 203 (2), pp. 293-301.

Perry T., et al., "Protection and Reversal of Excitotoxic Neuronal Damage by Glucagon-Like Peptide-1 and Exendin-4," *The Journal of Pharmacology and Experimental Therapeutics*, 2002, vol. 302 (3), pp. 881-888.

Perry T., et al., "The Glucagon-Like Peptides: A Double-Edged therapeutic Sword?," *Trends in Pharmacological Sciences*, 2003, vol. 24 (7), pp. 377-383.

Perry T.A., et al., "A New Alzheimer's Disease interventive Strategy: GLP-1," *Current Drug Targets*, 2004, vol. 5 (6), pp. 565-571.

Pillion D.J., et al., "Dodecylmaltoside-Mediated Nasal and Ocular Absorption of Lyspro-Insulin: Independence of Surfactant from Multimer Dissociation," *Pharmaceutical Research*, 1998, vol. 15(10), pp. 1637-1639.

Pinget M., et al., "Efficacy and safety of lixisenatide once daily versus placebo in type 2 diabetes insufficiently controlled on pioglitazone (GetGoal-P)," *Diabetes, Obesity and Metabolism*, 2013, vol. 15 (11), pp. 1000-1007.

Pohl M., et al., "Molecular Cloning of the Heloderma and Exendin-4 cDNAs in the Lizard," *The Journal of Biological Chemistry*, 1998, vol. 273 (16), pp. 9778-9784.

Porter D.W., et al., "Four Weeks Administration of Liraglutide Improves Memory and Learning As Well As Glycaemic Control in Mice with High Fat Dietary-induced Obesity and Insulin Resistance," *Diabetes, Obesity and Metabolism*, 2010, vol. 12 (10), pp. 891-899.

Pradier Let al. "Animal Models of Alzheimer's disease." *Demences (Dementias)*; eds. Duyckaerts C. and Pasquier F.; publisher Doin; 165-170 (Sep. 10, 2002; available Aug. 27, 2002).

Prandini N., "Methods of Measuring Gallbladder Motor Functions—the Need for Standardization: Scintigraphy," *Digestive and Liver Disease*, 2003, vol. 35 (Suppl 3), pp. S62-66.

"Preferable." Merriam-Webster.com. Merriam-Webster, n.d. Web. Sep. 7, 2015. <http://www.merriamwebster.com/dictionary/preferable>.

Pugeat M., et al., "Insulin Resistance, Polycystic Ovary Syndrome and Metformin," *Drugs*, 1999, vol. 58 (Suppl 1), pp. 41-46.

Quianzon C.L., et al., "Lixisenatide-Once Daily Glucagon-Like Peptide-1 Receptor Agonist in the Management of Type 2 Diabetes," *US Endocrinology*, 2011, vol. 7 (2), pp. 104-109.

Raccach D., et al., "When Basal Insulin therapy in Type 2 Diabetes Mellitus is not Enough—What Next?," *Diabetes/Metabolism Research and Reviews*, 2007, vol. 23 (4), pp. 257-264.

Raju R.P., et al., "Optimum Palliation of inoperable Hilar Cholangiocarcinoma: Comparative Assessment of the Efficacy of Plastic and Self-Expanding Metal Stents," *Digestive Diseases and Sciences*, 2011, vol. 56, pp. 1557-1564.

Ramos B., et al., "Early Neuropathology of Somatostatin/NPY GABAergic Cells in the Hippocampus of a PstAPP Transgenic Model of Alzheimer's Disease," *Neurobiology of Aging*, 2006, vol. 27 (11), pp. 1658-1672.

Rao A.D., et al., "Is the Combination of Sulfonylureas and Metformin Associated with an increased Risk of Cardiovascular Disease or all-Cause Mortality? A Meta-Analysis of Observational Studies," *Diabetes Care*, 2008, vol. 31 (8), pp. 1672-1678.

Ratner R.E., et al., "A Dose-Finding Study of the New GLP-1 Agonist Ave0010 in Type 2 Diabetes Insufficiently Controlled with Metformin," *Diabetes*, 2008, vol. 57 (Suppl 1), pp. A129. Abstract # 433-P.

Ratner R.E., et al., "Dose-Dependent Effects of the Once-Daily GLP-1 Receptor Agonist Lixisenatide in Patients with Type 2 Diabetes Inadequately Controlled with Metformin: A Randomized Double-Blind, Placebo-Controlled Trial," *Diabetic Medicine*, 2010, vol. 27 (9), pp. 1024-1032.

Ratner R.E., et al., "Post-Meal Pharmacodynamics Profile of Ave0010, A Once-Daily GLP-1 Receptor Agonist, in Patients with Type 2 Diabetes Inadequately Controlled on Metformin," *Diabetologia*, 2009, vol. 52 (Suppl 1), pp. S60. Abstract 131.

Ratner R.E., et al., "A Dose-Finding Study of the New GLP-1 Agonist AVE0010 in Type 2 Diabetes Insufficiently Controlled with Metformin," *Diabetes*, 68th Annual Meeting of the American Diabetes Association, San Francisco, Jun. 6-10, 2008, vol. 57 (Suppl 1), p. A129. Poster.

Request for "Type C" Meeting letter sent by Michael Lutz addressed to Mary Parks, dated Apr. 21, 2006, pp. 1-10.

Richter, von Margret, "Oldtimer as Newcomer" *Pharmazie*, pp. 1-9; <http://www.pharmazeutische-zeitung.de/pza/2002-12/pharm1.htm> (Feb. 2002).

Riddle M., et al., "Contributions of Basal and Postprandial Hyperglycemia over a Wide Range of a 1 C Levels before and after Treatment Intensification in Type 2 Diabetes," *Diabetes Care*, 2011, vol. 34, pp. 2508-2514.

Riddle M.C., et al., "Adding Once-Daily Lixisenatide for Type 2 Diabetes Inadequately Controlled by Established Basal Insulin: A 24-Week, Randomized, Placebo-Controlled Comparison (Getgoal-L)," *Diabetes Care*, 2013, vol. 36 (9), pp. 2489-2496.

Riddle M.C., et al., "Adding once-Daily Lixisenatide for Type 2 Diabetes inadequately Controlled with Newly initiated and Continuously Titrated Basal Insulin Glargine," *Diabetes Care*, 2013, pp. 2497-2503.

Ritzel U., et al., "A Synthetic Glucagon-Like Peptide-1 Analog with Improved Plasma Stability," *The Journal of Endocrinology*, 1998, vol. 159 (1), pp. 93-102.

Rohrmann C.A., "Differential Diagnosis of Pancreatic and Biliary Duct Diseases," *Diseases of the Abdomen and Pelvis Syllabus*, 1999, pp. 170-174.

Rosenstock et al., Abstract, "71st Scientific Sessions" <http://www.call4abstracts.com/ada/ada11d11b/index.php> 02:22:24 pp. 1-3, (Nov. 2011).

US 9,526,764 B2

Page 16

(56)

References Cited

OTHER PUBLICATIONS

Rosenstock J., et al., "Dose Range Effects of the New Once Daily GLP-1 Receptor Agonist Ave0010 Added to Metformin in Type 2 Diabetes," *Diabetologia*, 2008, vol. 51 (Suppl 1), pp. S66. Abstract 145.

Rosenstock J., et al., "Post-Meal Effects of Ave0010, A Once-Daily GLP-1 Receptor Agonist, in Type 2 Diabetes Inadequately Controlled on Metformin," *Diabetes*, 2009, vol. 58 (Suppl 1), pp. A151-A152. Abstract 564P.

Rosenstock J., et al., "Efficacy and Safety of Lixisenatide Once Daily vs Exenatide Twice Daily in Type 2 DM Inadequately Controlled on Metformin (GetGoal-X)," 71st Scientific Sessions, Nov. 2011. Poster.

Rubino A., et al., "Delayed initiation of Subcutaneous Insulin therapy after Failure of oral Glucose-Lowering Agents in Patients with Type 2 Diabetes: A Population-Based Analysis in the UK," *Diabetic Medicine*, 2007, vol. 24 (12), pp. 1412-1418.

Sampson H.A., et al., "Second Symposium on the Definition and Management of Anaphylaxis: Summary Report—Second National Institute of Allergy and Infectious Disease/Food Allergy and Anaphylaxis Network Symposium," *The Journal of Allergy and Clinical Immunology*, 2006, vol. 117 (2), pp. 391-397.

Sanger F., et al., "The Amide Groups of Insulin," *The Biochemical Journal*, 1955, vol. 59 (3), pp. 509-518.

Sanofi and Zealand Pharma Press Release (Evaluate), "Additional Positive Results from Global Phase III Program With Lixisenatide for Type 2 Diabetes", (Apr. 12, 2011) pp. 1-3.

Sanofi's Lantus Draft Prescribing Information/Package Insert: "NDA 21-081 DRAFT package insert" (Sponsor revision #5) Date of submission: Apr. 20, 2000; see http://www.drugbank.ca/system/fds_labels/DB00047.pdf 1265922812; pp. 1-14.

Sanofi Press Release entitled "FDA Accepts Sanofi's New Drug Application for Basal Insulin Toujeo®," dated Jul. 8, 2014, pp. 1-2. Sanofi Press Release; "Lyxumia (lixisenatide) in Combination with Basal insulin plus Oral Anti-Diabetics Significantly Reduced HbA1c and Post-Prandial Glucose"; Paris, France (Jun. 9, 2012) pp. 1-6.

Sanofi Press Release (Peron and Schaeffer), "Sanofi GetGoal Program on Lyxumia®, as an Add-on to Basal Insulin, Shows Significant Positive Phase III Results," Paris, France (May 31, 2011) pp. 1-2.

Sanofi Press release (Peron and Schaeffer); "Sanofi Reports Positive Results for Once-daily Lyxumia® (lixisenatide) in Combination with Lantus® (insulin glargine) in Type 2 Diabetes" Paris, France (Dec. 6, 2011) pp. 1-3.

Sanofi Press Release (Peron and Schaeffer), "Lyxumia® (lixisenatide) One-Step Regimen as Effective as Two-Step Regimen in Improving Glycemic Control in Type 2 Diabetes" Paris, France (Sep. 12, 2011) pp. 1-3.

Sanofi Press Release "Positive Results for Investigational Compound Lyxumia (Lixisenatide) Presented at American Diabetes Association's 71st Annual Scientific Sessions," (Jun. 24, 2011), pp. 1-5.

Sanofi Press Release (Sigurd), "Lixisenatide Significantly Reduces HbA1c Without Increasing Hypoglycemia in Patients Uncontrolled on Sulfonylureas", *Pressmeddelande* (Apr. 12, 2011) pp. 1-2.

Sanofi-aventis Press Release (Gabriel), "New Diabetes Compound AVE0010 Showed Clear Dose Response Results With Once-A-Day Injection in Phase IIb Study" Paris, France (Jun. 7, 2008) pp. 1-2. Sanofi-aventis Press Release, "Once Daily Lixisenatide (AVE 0010) Given as Monotherapy Successfully Meets Phase III Study Endpoints in Diabetes" Paris, France (Apr. 15, 2010) pp. 1-2.

Sanofi-aventis Press Release (Peron and Schaeffer), "Sanofi-aventis Announces Positive Top-line Lixisenatide Phase III Results" Paris, France (Feb. 2, 2011) pp. 1-2.

Sanofi Press Release entitled "Sanofi Receives FDA Approval of Once-Daily Basal Insulin Toujeo®," dated Feb. 26, 2015, pp. 1-4. Schapira A.H., "Causes of Neuronal Death in Parkinson's Disease," *Advances in Neurology*, 2001, vol. 86, pp. 155-162.

Schellenberger V., et al., "Attempts for Quantifying the S' Subsite Specificity of Serine Proteases, Selected Papers Presented at the 2nd International Meeting On the Molecular and Cellular Regulation of Enzyme Activity, *Advances in the Biosciences, Peptides and Proteases*," Recent Advances, 1987, vol. 65, pp. 159-166.

Schellenberger V., et al., "Protease-Catalyzed Kinetically Controlled Peptide Synthesis," *Angewante Chemie*, 1991, vol. 30 (11), pp. 1437-1449.

Schindowski K., et al., "Impact of Aging: Sporadic, and Genetic Risk Factors on Vulnerability to Apoptosis in Alzheimer's Disease," *Neuromolecular Medicine*, 2003, vol. 4 (3), pp. 161-178.

Schmitz C., et al., "Hippocampal Neuron Loss Exceeds Amyloid Plaque Load in a Transgenic Mouse Model of Alzheimer's Disease," *The American Journal of Pathology*, 2004, vol. 164 (4), pp. 1495-1502.

Schwartz G.P., et al., "A Superactive Insulin: [B10-Aspartic Acid]Insulin(Human)," *Proceedings of the National Academy of Sciences of the United States of America*, 1987, vol. 84 (18), pp. 6408-6411.

Search Report of the Indecopi for Patent Application in Peru No. 000643-2012/DIN, dated Jul. 23, 2015, pp. 1-2.

Search Report of the Intellectual Property Corporation of Malaysia for Malaysian Patent Application No. PI 2011006204; dated Sep. 15, 2015, pp. 1-3.

Secnik Boye K., et al., "Patient-Reported Outcomes in a Trial of Exenatide and Insulin Glargine for the Treatment of Type 2 Diabetes," *Health and Quality of Life Outcomes*, 2006, vol. 4 (80), pp. 1-8.

Seino Y., et al., "Report of the Committee on the Classification and Diagnostic Criteria of Diabetes Mellitus," *Journal of the Japan Diabetes Society*, 2010, vol. 53, pp. 450-467 (in Japanese) English summary also provided.

Seino Y., et al., "Randomized, Double-Blind, Placebo-Controlled Trial of the Once-Daily GLP-1 Receptor Agonist Lixisenatide in Asian Patients with Type 2 Diabetes Insufficiently Controlled on Basal Insulin with or without a Sulfonylurea (Getgoal-L-Asia)," *Diabetes, Obesity and Metabolism*, 2012, vol. 14 (10), pp. 910-917. Sharplin P., et al., "Improved Glycaemic Control by Switching from Insulin NPH to Insulin Glargine: A Retrospective Observational Study," *Cardiovascular Diabetology*, 2009, vol. 8 (3), pp. 1-8.

Sherer T.B., et al., "Subcutaneous Rotenone Exposure Causes Highly Selective Dopaminergic Degeneration and A-Synuclein Aggregation," *Experimental Neurology*, 2003, vol. 179, pp. 9-16.

Sluzky V., et al., "Kinetics of Insulin Aggregation in Aqueous Solutions Upon Agitation in the Presence of Hydrophobic Surfaces," *Proceedings of the National Academy of Sciences of the United States of America*, 1991, vol. 88 (21), pp. 9377-9381.

Smolka M.B., et al., "Optimization of the Isotope-Coded Affinity Tag-Labeling Procedure for Quantitative Proteome Analysis," *Analytical Biochemistry*, 2001, vol. 297 (1), pp. 25-31.

Sporn M.B., et al., "Chemoprevention of Cancer," *Carcinogenesis*, 2000, vol. 21 (3), pp. 535-530.

St. John Providence Health Center, "Preventing Obesity" <http://www.stjohnprovidence.org/HealthInfo/lib/swarticle.aspx?type=85&id=P07863>, Retrieved Aug. 22, 2013, pp. 1-2.

Starkova N.T., "Clinical Endocrinology," *Guide for physicians, Medicine*, 1991, pp. 192-262.

Stolk R.P., et al., "Insulin and Cognitive Function in an Elderly Population the Rotterdam Study," *Diabetes Care*, 1997, vol. 20 (5), pp. 792-795.

Summary of Product Characteristics Lyxumia 10 micrograms solution for injection, pp. 1-93, published Mar. 14, 2013.

Sundby F., "Separation and Characterization of Acid-induced Insulin Transformation Products by Paper Electrophoresis in 7 M Urea," *The Journal of Biological Chemistry*, 1962, vol. 237 (11), pp. 3406-3411.

Tanner C.M., et al., "Rotenone, Paraquat, and Parkinson's Disease," *Environmental Health Perspectives*, 2011, vol. 119 (6), pp. 866-872. Tempero M.A., "How I Treat Pancreatic Ductal Adenocarcinoma," *Current Clinical Issues, Journal of Oncology Practice*, 2008, vol. 4 (1), pp. 46-47.

Teramoto S., et al., "Exendin-4, a Glucagon-Like Peptide-1 Receptor Agonist, provides Neuroprotection in Mice Transient Focal

US 9,526,764 B2

Page 17

(56)

References Cited

OTHER PUBLICATIONS

Cerebral Ischemia,” *Journal of Cerebral Blood Flow and Metabolism*, 2011, vol. 31 (8), pp. 1696-1705.

Tessari P., et al., “Insulin in Methionine and Homocysteine Kinetics in Healthy Humans: Plasma Vs intracellular Models,” *American Journal of Physiology. Endocrinology and Metabolism*, 2005, vol. 288 (6), pp. E1270-E1276.

Tetich M., et al., “Neuroprotective Effects of (24R)-1,24-Dihydroxycholecalciferol in Human Neuroblastoma SH-SY5Y Cell Line,” *The Journal of Steroid Biochemistry and Molecular Biology*, 2004, vol. 89-90 (1-5), pp. 365-370.

The Expert Committee On The Diagnosis And Classification Of Diabetes Mellitus. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus Diabetes Care, Jan. 1998, 21:Supplement 1 S5-S19.

Thong K.Y., et al., “Safety, Efficacy and tolerability of Exenatide in Combination with Insulin in the Association of British Clinical Diabetologists Nationwide Exenatide Audit,” *Diabetes, Obesity and Metabolism*, 2011, vol. 13 (8), pp. 703-710.

Thurrow H., et al., “Stabilisation of Dissolved Proteins against Denaturation at Hydrophobic Interfaces,” *Diabetologia*, 1984, vol. 27 (2), pp. 212-218.

Toth M.L., et al., “Neurite Sprouting and Synapse Deterioration in the Aging Caenorhabditis Elegans Nervous System,” *The Journal of Neuroscience*, 2012, vol. 32 (26), pp. 8778-8790.

Translation of pp. 1109, 1116 and 1117 of “Clinical Effectiveness of Long-Term Administration of BAY g5421 (Acarbose) on Insulin-Treated Diabetes,” *Jpn. Pharmacol. Ther.*; 1996 vol. 24 No. 5: 1109-1129, pp. 1-4.

Translation of pp. 121 and 124 of Igaku to Yakugaku, “Utility of Voglibose Long-term Combined Therapy in Non-Insulin Dependent Diabetic Patients with Little Effective of Sulfonylureas,” 1999, vol. 42, No. 1: 121-129, pp. 1-3.

Translation of pp. 2346 and 2348 of Rinsho to Kenkyu, “Effectiveness of Combination Therapy Using Voglibose and Insulin in Patients with NIDDM,” 1997, vol. 74, No. 9: 2346-2352, pp. 1-3.

Translation of pp. 750, 753 and 754 of Igaku No Ayumi, “Incretin Receptors,” 2010, May, vol. 233; No. 9: 750-754, pp. 1-4.

Turner R.C., et al., “Glycemic Control with Diet, Sulfonylurea, Metformin, or Insulin in Patients with Type 2 Diabetes Mellitus: Progressive Requirement for Multiple therapies (UKPDS 49),” *JAMA*, 1999, vol. 281 (21), pp. 2005-2012.

Tyler-Cross R., et al., “Effects of Amino Acid Sequence, Buffers, and Ionic Strength on the Rate and Mechanism of Deamidation of Asparagine Residues in Small Peptides,” *The Journal of Biological Chemistry*, 1991, vol. 266 (33), pp. 22549-22556.

UK Prospective Diabetes Study (UKPDS) Group, “Effect of Intensive Blood-Glucose Control with Metformin on Complications in Overweight Patients with Type 2 Diabetes (UKPDS 34),” *Lancet*, 1998, vol. 352 (9131), pp. 854-865.

UK Prospective Diabetes Study (UKPDS) Group, “Intensive Blood-Glucose Control with Sulphonylureas or Insulin Compared with Conventional Treatment and Risk of Complications in Patients with Type 2 Diabetes (UKPDS 33),” *The Lancet*, 1998, vol. 352, pp. 837-853.

“Suspension” *Stedman’s Medical Dictionary*, 20th Edition, p. 1450 (Williams & Wilkins Co., Baltimore 1961).

“Suspension” *Taber’s Cyclopedic Medical Dictionary*, 19th Edition, p. 2097 (F.A. Davis Co., Philadelphia 2001).

Valle J., et al., “Cisplatin Plus Gemcitabine Versus Gemcitabine for Biliary Tract Cancer,” *The New England Journal of Medicine*, 2010, vol. 362 (14), pp. 1273-1281.

Van Delden, “Pancreas-Carcinoma, CT Assessment of Resectability,” *Radiology Department of the Academical Medical Centre*, Apr. 2006, pp. 1-12.

Varadarajan S., et al., “Review: Alzheimer’s Amyloid Beta-Peptide-Associated Free Radical Oxidative Stress and Neurotoxicity,” *Journal of Structural Biology*, 2000, vol. 130 (2-3), pp. 184-208.

Venezia V., et al., “Apoptotic Cell Death and Amyloid Precursor Protein Signaling in Neuroblastoma SH-SY5Y Cells,” *Annals of the New York Academy of Sciences*, 2004, vol. 1030, pp. 339-347.

Victoza® ANNEX I—Summary of product characteristics. First published 2009, pp. 1-32.

Victoza Press Release, “Diabetes drugs show promise in Alzheimer’s” published Jan. 17, 2011, pp. 1-2.

Victoza® Product information—European Medicines Agency, first published Aug. 7, 2009, pp. 1-2.

Volund A., et al., “In Vitro and in Vivo Potency of Insulin Analogues Designed for Clinical Use,” *Diabetic Medicine*, 1991, vol. 8 (9), pp. 839-847.

Vora J., et al., “Incretin-Based therapy in Combination with Basal Insulin: A Promising Tactic for the Treatment of Type 2 Diabetes,” *Diabetes & Metabolism*, 2013, vol. 39 (1), pp. 6-15.

Wafa W.S., et al., “Use of U-500 Regular Insulin in Type 2 Diabetes,” *Diabetes Care*, 2006, vol. 29 (9), pp. 2175-2176.

Wajchenberg B.L., “Clinical Approaches to Preserve Beta-Cell Function in Diabetes,” *Advances in Experimental Medicine and Biology*, 2010, vol. 654, pp. 515-535.

Wan Z., et al., “Enhancing the Activity of Insulin at the Receptor Interface: Crystal Structure and Photo-Cross-Linking of A8 Analogues,” *Biochemistry*, 2004, vol. 43 (51), pp. 16119-16133.

Wang L., et al., “Real-World Outcomes of US Employees with Type 2 Diabetes Mellitus Treated with Insulin Glargine or Neutral Protamine Hagedorn Insulin: A Comparative Retrospective Database Study,” *BMJ Open*, 2013, vol. 3 (4), pp. e002348 1-9.

Ward J.D., “Diabetic Neuropathy,” *British Medical Bulletin*, 1989, vol. 45 (1), pp. 111-126.

Watson G.S., et al., “Insulin increases CSF Abeta42 Levels in Normal Older Adults,” *Neurology*, 2003, vol. 60 (12), pp. 1899-1903.

Weiss M.A., et al., “Activities of Monomeric Insulin Analogs at Position A8 are Uncorrelated with their thermodynamic Stabilities,” *The Journal of Biological Chemistry*, 2001, vol. 276 (43), pp. 40018-40024.

Werner et al., “Abstract, Insulin Glargine U-100 Has a Favourable Time-Action Profile Compared to U-40 or NPH Insulin in Healthy, Normoglycaemic Dogs”, Poster 37th Annual Meeting of Endocrine Society of India, Tirupati, A.P., India, ESICON, 2007, p. 2.

Werner U., et al., “Pharmacological Profile of Lixisenatide: A New GLP-1 Receptor Agonist for the Treatment of Type 2 Diabetes,” *Regulatory Peptides*, 2010, vol. 164 (2-3), pp. 58-64.

White I.R., et al., “Randomized Clinical Trials with Added Rescue Medication: Some Approaches to their Analysis and interpretation,” *Statistics in medicine*, 2001, vol. 20 (20), pp. 2995-3008.

Whittingham J.L., et al., “Insulin At pH 2: Structural Analysis of the Conditions Promoting Insulin Fibre formation,” *Journal of Molecular Biology*, 2002, vol. 318 (2), pp. 479-490.

WHO BMI classification, accessed at URL apps.who.int/bmi/index.jsp?introPage=itrol_3.html, Sep. 9, 2013, one page.

WHO Drug Information vol. 22(2), list 99, p. 142 (lixisenatide) (Jul. 2008).

Widjaja A., et al., “UKPDS 20: Plasma Leptin, Obesity, and Plasma Insulin in Type 2 Diabetic Subjects,” *The Journal of Clinical Endocrinology and Metabolism*, 1997, vol. 82 (2), pp. 654-657.

Wiernsperger N.F., et al., “The Antihyperglycaemic Effect of Metformin: Therapeutic and Cellular Mechanisms,” *Drugs*, 1999, vol. 58 (Suppl 1), pp. 31-39.

Wirths O., et al., “Intraneuronal Abeta Accumulation Precedes Plaque formation in beta-Amyloid Precursor Protein and Presenilin-1 Double-Transgenic Mice,” *Neuroscience Letters*, 2001, vol. 306 (1-2), pp. 116-120.

Wirths O., et al., “Intraneuronal APP/A Beta Trafficking and Plaque formation in Beta-Amyloid Precursor Protein and Presenilin-1 Transgenic Mice,” *Brain Pathology*, 2002, vol. 12 (3), pp. 275-286.

Wirths O., et al., “Reelin in Plaques of Beta-Amyloid Precursor Protein and Presenilin-1 Double-Transgenic Mice,” *Neuroscience Letters*, 2001, vol. 316 (3), pp. 145-148.

Wollen K.A., “Alzheimer’s Disease: the Pros and Cons of Pharmaceutical, Nutritional, Botanical, and Stimulatory therapies, with a

(56)

References Cited

OTHER PUBLICATIONS

Discussion of Treatment Strategies from the Perspective of Patients and Practitioners,” *Alternative Medicine Review*, 2010, vol. 15 (3), pp. 223-244.

Workgroup on Hypoglycemia, American Diabetes Association, “Defining and Reporting Hypoglycemia in Diabetes: A Report from the American Diabetes Association Workgroup on Hypoglycemia,” *Diabetes Care*, 2005, vol. 28 (5), pp. 1245-1249.

Written Opinion of the ISA for International Application No. PCT/EP2011/058079, mailed Mar. 22, 2012, pp. 1-8.

Xie H., et al., “Characterization of Protein Impurities and Site-Specific Modifications Using Peptide Mapping with Liquid Chromatography and Data Independent Acquisition Mass Spectrometry,” *Analytical Chemistry*, 2009, vol. 81 (14), pp. 5699-5708.

Yki-Jarvinen H., et al., “Insulin Glargine or Nph Combined with Metformin in Type 2 Diabetes: The Lanmet Study,” *Diabetologia*, 2006, vol. 49 (3), pp. 442-451.

Yki-Jarvinen H., “Thiazolidinediones,” *The New England Journal of Medicine*, 2004, vol. 351 (11), pp. 1106-1118.

Yoon N.M., et al., “Exenatide Added to Insulin therapy: A Retrospective Review of Clinical Practice Over Two Years in an Academic Endocrinology Outpatient Setting,” *Clinical Therapeutics*, 2009, vol. 31 (7), pp. 1511-1523.

Yu Z.P., et al., “Effect of Zinc Sulphate and Zinc Methionine on Growth, Plasma Growth Hormone Concentration, Growth Hormone Receptor and Insulin-Like Growth Factor-I Gene Expression in Mice,” *Clinical and Experimental Pharmacology & Physiology*, 2005, vol. 32 (4), pp. 273-278.

Zealand Pharma Press Release entitled “Sanofi-Aventis finalize phase Iia clinical study with GLP-1 agonist for type 2 diabetes licensed from Zealand Pharma” dated Mar. 3, 2005, one page.

Ziemer D.C., et al., “Clinical Inertia Contributes to Poor Diabetes Control in a Primary Care Setting,” *The Diabetes Educator*, 2005, vol. 31 (4), pp. 564-571.

Ziessman H.A., et al., “Sincalide-Stimulated Cholescintigraphy: A Multicenter Investigation to Determine Optimal Infusion Methodology and Gallbladder Ejection Fraction Normal Values,” *Journal of Nuclear Medicine*, 2010, vol. 51 (2), pp. 277-281.

Zimmet P., et al., “Clinical Efficacy of Metformin Against Insulin Resistance Parameters: Sinking the Iceberg,” *Review Article, Drugs*, 1999, vol. 58, (Suppl 1), pp. 21-28.

Zinman B., et al., “Efficacy and Safety of the Human Glucagon-Like Peptide-1 Analog Liraglutide in Combination with Metformin and Thiazolidinedione in Patients with Type 2 Diabetes (LEAD-4 Met+TZD),” *Diabetes Care*, 2009, vol. 32 (7), pp. 1224-1230.

Zinman B., “The Physiologic Replacement of Insulin an Elusive Goal,” *The New England Journal of Medicine*, 1989, vol. 321 (6), pp. 363-370.

* cited by examiner




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5a




Fig. 5b




Fig. 6




Fig. 7




Fig. 8




Fig. 9

Lantus dose	AVE10 dose		
	Mix A	Mix B	Mix C
10	6.7		
12	8.0		
14	9.3		
16	10.7		
18	12.0		
20	13.3		
22	14.7	8.8	
24	16.0	9.6	
26	17.3	10.4	
28	18.7	11.2	
30	20.0	12.0	
32		12.8	
34		13.6	
36		14.4	
38		15.2	
40		16.0	
42		16.8	10.5
44		17.6	11.0
46		18.4	11.5
48		19.2	12.0
50		20.0	12.5
52			13.0
54			13.5
56			14.0
58			14.5
60			15.0
62			15.5
64			16.0
66			16.5
68			17.0
70			17.5
72			18.0
74			18.5
76			19.0
78			19.5
80			20.0

Fig. 10

US 9,526,764 B2

1

**COMBINATION OF AN INSULIN AND A
GLP-1-AGONIST**

The invention relates to a medicament comprising at least one insulin and at least one GLP-1 receptor agonist, referred to below as GLP-1 agonist, the medicament being formulated and/or compounded in such a way that it comprises the insulin and the GLP-1 agonist each in a predetermined amount and can be administered in a dose adapted to the individual requirement of a patient.

The invention relates more particularly to a medicament comprising a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, which each comprise at least one insulin and at least one GLP-1 agonist and contain the at least one insulin and/or the at least one GLP-1 agonist in different fractions relative to the total weight of the composition.

More particularly the present invention relates to a medicament which comprises a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, the first pharmaceutical composition comprising at least one insulin, and the second pharmaceutical composition comprising at least one insulin and at least one GLP-1 agonist, and the at least one further pharmaceutical composition comprising at least one insulin and at least one further active compound.

Around 250 million people worldwide suffer from diabetes mellitus. For the type 1 diabetics among them, replacement of the deficient endocrine insulin secretion is the only possible therapy at present. Those affected are dependent on insulin injections for life, usually several times a day. Type 2 diabetes contrasts with type 1 diabetes in that there is not always a deficiency of insulin, but in a large number of cases, especially at the advanced stage, treatment with insulin, where appropriate in combination with an oral antidiabetic, is considered the most advantageous form of therapy.

In healthy individuals, insulin release by the pancreas is strictly coupled to the blood glucose concentration. Elevated blood glucose levels like those occurring after meals are rapidly compensated by a corresponding rise in insulin secretion. In the fasting state, the plasma insulin level falls to a baseline value which is sufficient to ensure a continuous supply of glucose to insulin-sensitive organs and tissues and to keep hepatic glucose production low during the night. The replacement of the endogenous insulin secretion by exogenous, usually subcutaneous administration of insulin does not in general come close to the above-described quality of the physiological regulation of blood glucose. Frequently there are instances of blood glucose being thrown off-track, either upwardly or downwardly, and in their most severe forms these may be life-threatening. In addition, however, blood glucose levels which are elevated over years, without initial symptoms, constitute a considerable health risk. The large-scale DCCT study in the USA (The Diabetes Control and Complications Trial Research Group (1993) N. Engl. J. Med. 329, 977-986) showed unambiguously that chronically elevated blood glucose levels are responsible for the development of late diabetic complications. Late diabetic complications are micro- and macrovascular damage which is manifested in certain circumstances as retinopathy, nephropathy or neuropathy and leads to blindness, renal failure, and loss of extremities, and, in addition, is associated with an increased risk of cardiovascular disorders. From this it can be inferred that an improved therapy of diabetes must be aimed primarily at keeping blood glucose as closely as

2

possible within the physiological range. According to the concept of intensified insulin therapy, this is to be achieved by means of injections, several times a day, of fast-acting and slow-acting insulin preparations. Fast-acting formulations are given at mealtimes in order to compensate the postprandial rise in blood glucose. Slow-acting basal insulins are intended to ensure the basic supply of insulin especially during the night, without leading to hypoglycemia.

Insulin is a polypeptide composed of 51 amino acids which are divided over 2 amino acid chains: the A chain, with 21 amino acids, and the B chain, with 30 amino acids. The chains are linked together by 2 disulfide bridges. Insulin preparations have been employed for many years for diabetes therapy. Such preparations use not only naturally occurring insulins but also, more recently, insulin derivatives and insulin analogs.

Insulin analogs are analogs of naturally occurring insulins, namely human insulin or animal insulins, which differ by replacement of at least one naturally occurring amino acid residue by other amino acids and/or addition/deletion of at least one amino acid residue from the corresponding, otherwise identical, naturally occurring insulin. The amino acids in question may also be amino acids which do not occur naturally.

Insulin derivatives are derivatives of naturally occurring insulin or of an insulin analog which are obtained by chemical modification. The chemical modification may consist, for example, in the addition of one or more defined chemical groups and to one or more amino acids. Generally speaking, the activity of insulin derivatives and insulin analogs is somewhat altered as compared with human insulin.

Insulin analogs with an accelerated onset of action are described in EP 0 214 826, EP 0 375 437, and EP 0 678 522. EP 0 214 826 relates, among other things, to replacements of B27 and B28. EP 0 678 522 describes insulin analogs which have different amino acids in position B29, preferably proline, but not glutamic acid. EP 0 375 437 encompasses insulin analogs with lysine or arginine in B28, which may optionally also be modified in B3 and/or A21. An accelerated activity is also exhibited by the insulin analogs described in EP-A-0 885 961.

EP 0 419 504 discloses insulin analogs which are protected from chemical modifications by modification of asparagine in B3 and of at least one further amino acid in positions A5, A15, A18 or A21.

WO 92/00321 describes insulin analogs in which at least one amino acid in positions B1-B6 has been replaced by lysine or arginine. Such insulins, according to WO 92/00321, have an extended activity. A delayed activity is also exhibited by insulin analogs described in EP-A 0 368 187 and by the insulin analogs described in German patent applications 10 2008 003 568.8 and 10 2008 003 566.1.

The insulin preparations of naturally occurring insulins for insulin replacement that are on the market differ in the origin of the insulin (e.g., bovine, porcine, human insulin) and also in their composition, whereby the profile of action can be influenced (onset of action and duration of action). By combining different insulin products it is possible to obtain a wide variety of profiles of action and to set blood sugar levels which are as close as possible to physiological. Recombinant DNA technology nowadays allows the production of such modified insulins. These include insulin glargine (Gly(A21)-Arg(B31)-Arg(B32) human insulin), with an extended duration of action. Insulin glargine is injected as an acidic, clear solution and, on account of its

US 9,526,764 B2

3

solution properties in the physiological pH range of the subcutaneous tissue, is precipitated as a stable hexamer associate. Insulin glargine is injected once daily and is notable over other long-activity insulins for its flat serum profile and the associated reduction in the risk of nocturnal hypoglycemia (Schubert-Zsilavec et al., 2:125-130 (2001)).

The specific preparation of insulin glargine that leads to a prolonged duration of action is characterized by a clear solution with an acidic pH.

Glucagon-like peptide 1 (GLP-1) is an endocrine hormone which increases the insulin response following oral intake of glucose or fat. GLP-1 generally regulates the concentrations of glucagons, slows down gastric emptying, stimulates the biosynthesis of (Pro-)insulin, increases the sensitivity toward insulin, and stimulates the insulin-independent biosynthesis of glycogen (Hoist (1999), *Curr. Med. Chem.* 6:1005, Nauck et al. (1997) *Exp Clin Endocrinol Diabetes* 105: 187, Lopez-Delgado et al. (1998) *Endocrinology* 139:2811).

Human GLP-1 has 37 amino acid residues (Heinrich et al., *Endocrinol.* 115:2176 (1984), Uttenthal et al., *J Clin Endocrinol Metabol* (1985) 61:472). Active fragments of GLP-1 include GLP-1(7-36) amide and GLP-1(7-37).

Exendins are another group of peptides which are able to lower blood glucose concentrations. Exendins have a certain similarity in sequence to GLP-1(7-36) (53%, Goke et al. *J. Biol Chem* 268, 19650-55). Exendin-3 and exendin-4 stimulate an increase in cellular cAMP production in acinar cells of the guinea pig pancreas by interaction with exendin receptors (Raufman, 1996, *Reg. Peptides* 61:1-18). In con to exendin-4, exendin-3 produces an increase in amylase release in acinar cells of the pancreas.

Exendin-3, exendin-4, and exendin agonists have been proposed for the treatment of diabetes mellitus and the prevention of hyperglycemia; they reduce gastric motility and gastric emptying (U.S. Pat. No. 5,424,286 and WO98/05351).

Exendin analogs may be characterized by amino acid replacements and/or C-terminal truncation of the natural exendin-4 sequence. Exendin analogs of this kind are described in WO 99/07404, WO 99/25727, WO 99/25728.

Combinations of insulin and GLP-1 are known from WO 2004/005342 for the treatment of diabetes.

In clinical practice the amount of insulin to be administered is adjusted to the individual requirement of the individual diabetes patients. Individual patients generally need different amounts of insulin and/or GLP-1 agonist. Typically the predetermined dose is administered by administering a defined amount of a composition having a defined concentration. A result of this is that a composition which comprises insulin and GLP-1 at the same time allows the administration of only one particular proportion of insulin and GLP-1. This means that only one of the two amounts of insulin and GLP-1 can be adapted optimally to the requirement of the patients. Since in practice the correct adjustment of the amount of insulin administered is essential, it is assumed that, when a particular proportion of insulin to GLP-1 is administered, the GLP-1 agonist is either underdosed or overdosed and is correct by chance at best.

There are various systems known for injecting a combination of active compounds. The active compounds may be formulated in a composition and provided in a device, as for example in a prefilled syringe. A system of this kind does allow the dosing of the combination, but only in a fixed portion of the active compounds, as is present in the composition. As set out therein, this is a disadvantage for the

4

combination of an insulin with a GLP-1 agonist, since different amounts of the insulins and of the GLP-1 agonist have to be administered, according to the therapeutic requirement.

It is also possible for two active compounds to be administered in two separate formulations, each comprising one of the two active compounds, which are injected independently of one another each with one device (e.g., prefilled syringes). In the case of an injection therapy such as the injection of insulin, for example, patient compliance is a key prerequisite for the success of the therapy. Generally speaking, in the case of an injection therapy, pain, needle-phobia, and the carrying facility for the injection apparatus are a problem, which can lead to reduced compliance. If the patient is to use two separate devices for injection, then these problems multiply.

A single device for the administration of insulin and a GLP-1 agonist is advantageous over the use of two separate devices for administering insulin and a GLP-1 agonist as far as the patient/user is concerned. Moreover, the use of only one device rather than two devices may reduce the number of steps which the patient/user must carry out, which lowers the frequency of errors in use. This reduces the risk of unwanted side effects.

U.S. Pat. No. 4,689,042, U.S. Pat. No. 5,478,323, U.S. Pat. No. 5,253,785, and WO 01/02039 describe devices for the simultaneous administration of two injectable product to a patient. These devices comprise two containers each containing one composition. In these devices the two compositions are injected via a needle. This does make it possible to overcome the disadvantages produced by the use of two separate devices. As a result of the mixing process, there is a dilution in the concentrations of the two active compounds. This may impact adversely on the pharmacokinetics.

The pharmacokinetics of insulin, particularly of insulin glargine, is influenced by the dilution of the insulin in the administered composition. In order to ensure reliable activity of a particular dose of insulin, therefore, the concentration of insulin ought to be kept constant as far as possible. Dosing ought to take place essentially via the volume of the insulin composition administered. This is also true for the administration of a combination of insulin and a GLP-1 agonist. When a combination of insulin and a GLP-1 agonist is administered, this proviso can only be met if both substances are dosed in a fixed proportion to one another in one composition. Where both substances are provided in separate compositions and are mixed for injection in a suitable device (e.g., from WO 01/02039), then a constant concentration of insulin can be realized only if the insulin composition is not substantially diluted by the composition of the GLP-1 agonist. This imposes limits on the possibility of independent dosing of insulin and of the GLP-1 agonist.

One conceivable solution would be to provide the GLP-1 agonist in such a high concentration that the dosed addition of the GLP-1 agonist produces not significant dilution of the insulin composition (e.g., not more than 10%). Polypeptides such as insulins (e.g., insulin glargine, Lantus®) or GLP-1 agonists cannot be concentrated ad infinitum. First, the solubility of proteins is limited, and high concentrations of protein may alter the flow characteristics of the solution. The most important problem for the use of solutions with a high concentration of active compound is the dosing accuracy. At high concentrations it would be necessary to administer small volumes or to carry out dosing into a different solution. There are devices known for the precise dosing of small or very small volumes. However, such devices are expensive

US 9,526,764 B2

5

and, on the basis of their operation, are intended only for use by trained personnel, as in the laboratory, for example. Since, generally speaking, patients inject themselves with insulins and/or GLP-1 agonists, the use of such devices for administering insulins and/or GLP-1 agonists is ruled out. The devices that are described, for example, in U.S. Pat. No. 4,689,042, U.S. Pat. No. 5,478,323, U.S. Pat. No. 5,253,785, and WO 01/02039, which allow the patients to inject themselves with active compound solutions, are unsuitable for the dosing of small and very small volumes.

The problems which arise with the injection of a combination of an insulin and a GLP-1 agonist are as follows

the proportion of the active compounds must be variable; the pharmacokinetics of at least one of the active compounds (the insulin) is influenced by the concentration/dilution;

the pharmacokinetics of at least one other active compound (the GLP-1 agonist) is not, or not substantially, influenced by the concentration/dilution.

It was an object of the present invention, therefore, to provide a medicament which at least partly overcomes the above-described disadvantages of the prior art. A further intention is that there should as far as possible be only one administration per day.

It has been found, surprisingly, that the combination of an insulin with a GLP-1 agonist exhibits synergistic effects in the regulation of blood glucose in the postprandial and postabsorptive phases as compared with the use of insulin or the GLP-1 agonist alone;

Higher activity on the basis of the combination of the complementary activities on the fasting and postprandial glucose levels, which complement one another (examples 2 and 3). The combination exhibits a lowering in postprandial glucose concentration (i.e., improved glucose tolerance) like a GLP-1 agonist alone, and additionally exhibits the postabsorptive lowering of glucose like an insulin (example 9).

Reduction in the risk of hypoglycemia (examples 2-4).

Improved adaptation of the blood glucose concentration to normoglycemic levels (example 8).

Improved glucose tolerance and lowering of postabsorptive glucose concentrations (example 9).

The synergistic activities of the combination on the glucose concentration are observed in a GLP-1 agonist concentration range of one order of magnitude (factor 10). (Example 6 compared with examples 4 and 2). Only in the case of relatively small GLP-1 doses and/or relatively large proportions of insulin to GLP-1 are the activities of insulin predominant.

Maintains the function of the β -cells (example 10).

Weight loss/reduction in weight gain.

All examples show that GLP-1 agonists and insulins exhibit no adverse interactions.

As a result of the activities on the fasting, postprandial, and postabsorptive blood glucose concentrations, it becomes possible to reduce the number of administrations of the combination to once daily.

The invention provides a medicament comprising at least one insulin and at least one GLP-1 agonist, the medicament being formulated and/or compounded in such a way that it comprises the insulin and the GLP-1 agonist each in a predetermined amount and can be administered in a dose adapted to the individual requirement of a patient.

The medicament of the invention is used in particular for treating patients with diabetes mellitus, more particularly patients with type 1 or type 2 diabetes.

6

The medicament of the invention allows the blood glucose concentration to be adapted more effectively to normoglycemic levels in the case of patients with diabetes, more particularly type 1 or type 2 diabetes.

The medicament is used preferably to adjust the fasting, postprandial and/or postabsorptive blood glucose concentration of patients with diabetes, more particularly patients with type 1 or type 2 diabetes. More preferably the medicament of the invention is used to adjust the postprandial and/or postabsorptive blood glucose concentration of patients with diabetes, more particularly patients with type 1 or type 2 diabetes. Adjustment in this context means that normoglycemic blood glucose concentrations are substantially achieved or at least an approximation thereto is obtained. By normoglycemic levels are meant more particularly blood glucose concentrations in the normal range (breadth of fluctuation 60-140 mg/dl, corresponding to 3.3 to 7.8 mmol/l). This range of fluctuation encompasses blood glucose concentrations under fasting conditions, postprandial conditions, and postabsorptive conditions.

Postprandial and postabsorptive are terms familiar to the person skilled in the field of diabetology. Postprandial is used herein to refer more particularly to the phase after a meal and/or after glucose loading in an experiment. This phase is characterized more particularly in a healthy individual by an increase and fall again in the concentration of glucose in the blood. Postabsorptive, or postabsorptive phase, is used herein to refer more particularly to the phase which follows the postprandial phase. The postprandial phase typically ends up to 4 h after the meal and/or glucose loading. The postabsorptive phase lasts typically for up to 8 to 16 h.

The medicament of the invention is also used preferably for improving glucose tolerance in the treatment of a patient with diabetes, more particularly with a type 1 or type 2 diabetes. Improving the glucose tolerance means that the medicament of the invention lowers the postprandial blood glucose concentration. Improving the glucose tolerance is also taken to mean that the medicament of the invention lowers the postabsorptive blood glucose concentration. Lowering means more particularly that the blood glucose concentration substantially reaches normoglycemic values or at least is approximated thereto.

The medicament of the invention is able to lower the risk of hypoglycemia, which may occur, for example, in the postabsorptive phase. The medicament of the invention is used preferably for preventing hypoglycemia in the treatment of a patient with diabetes, more particularly with a type 1 or type 2 diabetes, it being possible for the hypoglycemia to occur more particular in the postabsorptive phase.

The medicament of the invention may maintain the function of the pancreatic β -cells. The medicament of the invention is used preferably for preventing a loss of function of the pancreatic β -cells in a patient with diabetes, more particularly with a type 1 or type 2 diabetes. The loss of function of the β -cells may be caused more particularly by apoptosis.

Furthermore, the medicament of the invention may bring about a loss in weight and/or prevent a gain in weight in patients with diabetes, more particularly type I or II. In diabetes patients, especially those of type 2, weight gain and excessive weight are frequent problems. Accordingly, administering the medicament of the invention may support a therapy for the treatment of excessive weight.

It will be appreciated that the medicament of the invention can be used in order to treat more than one of the preferred indications described therein in a patient with diabetes, more

US 9,526,764 B2

7

particularly with a type 1 or 2 diabetes. Accordingly the present invention encompasses not only the individual preferred indications but also arbitrary combinations of the indications. The medicament of the invention can therefore be used to treat one or more of the herein-described indications in patients with diabetes, more particularly of patients with type 1 or type 2 diabetes, for the purpose, for example, of adjusting the fasting, postprandial and/or post-absorptive blood glucose concentration, for improving glucose tolerance, for preventing hypoglycemia, for preventing a loss of function of the pancreatic β -cells, for weight loss and/or for preventing weight gain. Preference is given to the adjustment of fasting, postprandial and/or postabsorptive blood glucose concentration, the improvement of glucose tolerance and/or the prevention of hypoglycemia.

The medicament of the invention can also be used for producing a medicinal product for treating one or more of the herein-described indications, as, for example, for adjusting the fasting, postprandial and/or postabsorptive blood glucose concentration, for improving glucose tolerance, for preventing hypoglycemia, for preventing a loss of function of the pancreatic β -cells, for weight loss and/or for preventing weight gain.

The at least one insulin and the at least one GLP-1 agonist may also be used for producing a medicinal product for treating one or more of the herein-described indications, as for example for adjusting the fasting, postprandial and/or postabsorptive blood glucose concentration, for improving glucose tolerance, for preventing hypoglycemia, for preventing a loss of function of the pancreatic β -cells, for weight loss and/or for preventing weight gain.

The at least one GLP-1 agonist and the at least one insulin may be provided together in one pharmaceutical composition. In this case a first and a second composition and, optionally, at least one further pharmaceutical composition are provided, each comprising the insulin and the GLP-1 agonist. Accordingly the invention provides a medicament comprising a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, each comprising at least one insulin and at least one GLP-1 agonist, and containing the at least one insulin and/or the at least one GLP-1 agonist in different weight fractions relative to the total weight of the composition.

In the present specification "optionally, at least one further pharmaceutical composition" means that the medicament of the invention, in addition to the first and second pharmaceutical compositions, may comprise at least one further pharmaceutical composition. Hence the medicament of the invention may comprise, for example, 3, 4, 5, 6, 7, 8, 9, 10 or more pharmaceutical compositions of the invention.

Preferred medicaments are those which comprise a first and a second pharmaceutical composition of the invention.

Likewise preferred are medicaments which comprise a first, a second, and a third pharmaceutical composition of the invention.

Likewise preferred are medicaments which comprise a first, a second, a third, and a fourth pharmaceutical composition of the invention.

Likewise preferred are medicaments which comprise a first, a second, a third, a fourth, and a fifth pharmaceutical composition.

The weight fractions of the at least one insulin and of the at least one GLP-1 agonist may be selected in the first pharmaceutical composition, the second pharmaceutical composition, and, where used, the at least one further pharmaceutical composition in such a way that the pharma-

8

ceutical compositions contain different proportions of insulin to GLP-1 agonist, based on the weight fraction.

In this case the first composition may contain the smallest proportion and the second composition the next-greater proportion. Where at least one further composition is present, it may contain the next-greater proportion. Where a further composition is present as well, it may contain the next-greater proportion in turn. The compositions may therefore contain proportions of insulin to GLP-1 agonist, based on the weight fraction, that increase from the first to the second and, where used, further compositions.

The weight fraction of one of the two active compounds, i.e., of the at least one insulin or of the at least one GLP-1 agonist, in the first pharmaceutical composition, the second pharmaceutical composition, and, where used, the at least one further pharmaceutical composition is preferably selected in each case such that the predetermined dose of this active compound can be administered by administering a defined volume of the first, second and/or at least one further composition. With particular preference this active compound is the at least one insulin.

The weight fraction of the other of the two active compounds, i.e., of the at least one insulin or of the at least one GLP-1 agonist, in the first pharmaceutical composition, the second pharmaceutical composition, and, where used, the at least one further pharmaceutical composition is preferably selected such that the proportions of insulin to GLP-1 agonist, based on the weight fraction, increase from the first to the second and, where used, further compositions. With particular preference this active compound is the at least one GLP-1 agonist.

Furthermore, the weight fraction of the other of the two active compounds in the pharmaceutical compositions is determined such that one of the pharmaceutical compositions can be selected in such a way that the dose of the first of the two active compounds that is to be administered and the dose of the second active compound that is to be administered are given in a defined volume. Hence a pharmaceutical composition is selected which contains the desired proportion.

Theoretically it would be possible to provide a pharmaceutical composition for each individual therapeutically desired proportion of the weight fractions of the at least one insulin to the at least one GLP-1 agonist, in order to obtain an optimum dosage, tailored to requirements, for both active compounds for every patient.

In the present invention, a particular number of pharmaceutical compositions is sufficient in order to cover the dosages needed in practice for the two active compounds. For each patient a defined dosage range is defined within a therapeutically rational interval for each of the two active compounds. The dose to be administered ought hereby to fluctuate essentially within this dosage range for a particular patient, without any overdosing or underdosing.

Surprisingly it has been found that the synergistic effects of the combination of at least one insulin and at least one GLP-1 agonist on the concentration of glucose in the blood plasma occur in a concentration range of the GLP-1 agonist of one order of magnitude (factor 10). Since it is primarily the amount of insulin that must be adapted and precisely dosed to the individual patient, the synergistic concentration range of the GLP-1 agonist allows a pharmaceutical composition of the invention that contains a defined proportion of at least one insulin to the at least one GLP-1 agonist to cover a therapeutic range of insulin doses simultaneously with the associated, synergistic amount of GLP-1 agonist. The proportion can be selected such that every desired

US 9,526,764 B2

9

insulin dose has its corresponding dose of the at least one GLP-1 agonist, which is situated within the desired range, e.g., the synergistic range. As set out earlier on above, the proportions of the first, second, and, where used, at least one further composition of the medicament may also be chosen such that the proportions increase from the first to the second and, where used, the at least one further composition. If the dose of the GLP-1 agonist at the desired insulin dose of a composition (e.g., of the first composition) is outside (generally above) the desired dosage range of the GLP-1 agonist, then the next composition (e.g., the second composition) or a further composition with a greater proportion of the at least one insulin to the at least one GLP-1 agonist, is selected for use, in which the amount of the GLP-1 agonist at the desired insulin dose lies within the desired range. The proportions of the first, second, and, where used, at least one further composition of the medicament may further be chosen such that the ranges of the insulin dosages which correspond to the desired dosages of the at least one GLP-1 agonist border one another and/or overlap one another. Preferably the ranges overlap. Overlapping means more particularly that it is possible to select at least two compositions which, at the desired dose of the at least one insulin, each contain an amount of the at least one GLP-1 agonist which lies within the desired dosage range.

For example, three compositions are sufficient to adjust the dose of the a least one insulin for an individual patient to a level selected from the range from 15 to 80 units of insulin and at the same time to dose the GLP-1 agonist with an amount within the range from 10 to 20 μg (see example 11).

It is also possible to provide a medicament of the invention in which the proportion is selected such that for each desired dosage of the GLP-1 agonist there is a corresponding dosage of the at least one insulin which lies within the desired range, e.g., the synergistic range. The proportions of the first, second, and, where used, at least one further composition of the medicament may also be chosen such that the ranges of the dosages of the GLP-1 agonist that correspond to the desired dosages of the at least one insulin border one another and/or overlap one another. Preferably the ranges overlap. Overlapping in this context means more particularly that it is possible to select at least two compositions which, at the desired dosage of the at least one GLP-1 agonist, each contain an amount of the at least one insulin that lies within the desired dosage range.

Preferably the medicament of the invention contains not more than 10 pharmaceutical compositions as defined above, more preferably not more than 5, not more than 4, not more than 3 or 2 pharmaceutical compositions.

The compositions of the invention may contain the at least one insulin in identical or different weight fractions. For example, at least two of the compositions of the invention may contain the at least one insulin in a substantially identical weight fraction.

It is preferred for the first, second, and, where used, further compositions to contain the at least one insulin in a substantially identical weight fraction and the at least one GLP-1 agonist in different weight fractions.

The compositions of the invention may contain the at least one GLP-1 agonist in identical or different weight fractions. For example, at least two of the compositions of the invention may contain the at least one GLP-1 agonist in a substantially identical weight fraction.

It is also preferred for the first, second, and, where used, further compositions to contain the at least one GLP-1

10

agonist in a substantially identical weight fraction and the at least one insulin in different weight fractions.

Besides the first, second, and, where used, at least one further composition, the medicament of the invention may comprise at least one further pharmaceutical composition which contains either at least one insulin or at least one GLP-1 agonist. The medicament of the invention may also comprise at least one further pharmaceutical composition which contains at least one insulin and at least one GLP-1 agonist in a proportion of the weight fractions which is like the herein-described first, second or, where used, further pharmaceutical composition.

The invention further provides a medicament comprising a first pharmaceutical composition and a second pharmaceutical composition, the first pharmaceutical composition comprising at least one insulin and the second pharmaceutical composition comprising at least one GLP-1 agonist, the medicament being formulated and/or compounded for the independent administration of the first and second pharmaceutical compositions.

Example 12 shows how a combination of two or more active compounds can be formulated such that when two or more compositions are combined, both active compounds can be administered in any desired amounts and in any desired proportions to one another. This takes account of the fact that at least one of the active compounds must not be diluted as a result of the combining (e.g., through mixing immediately prior to administration).

The present invention provides a medicament which comprises a first active compound and a second active compound, and, optionally, at least one further active compound, these active compounds being provided in a first, a second, and, optionally, at least one further composition. The first active compound is present in all of the compositions. The second active compound is present in the second formulation, and the at least one further active compound, where used, is present in the optionally at least one further composition. Hence the second and each further composition comprise the first active compound in combination with another active compound.

The present invention thus further provides a medicament which comprises a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, the first pharmaceutical composition comprising at least one first active compound, and the second pharmaceutical composition comprising at least one first active compound and at least one second active compound, and the at least one further pharmaceutical composition comprising at least one first and at least one further active compound. The active compounds here may be any desired active compounds.

The first composition preferably comprises as active compound only the at least one first active compound.

The first, second, and, where used, at least one further compositions may comprise the first active compound in a substantially identical weight fraction or in different weight fractions relative to the total weight of the composition.

It is preferred for the first pharmaceutical composition, the second pharmaceutical composition, and, where used, the at least one further pharmaceutical composition to comprise the first active compound in substantially equal weight fractions relative to the total weight of the composition. By this means it is possible to ensure that any desired proportion of the first and second composition and, where appropriate, any desired proportion of the first and at least one further composition can be used, the dosing of the first active compound taking place via the total amount of the compo-

US 9,526,764 B2

11

sitions administered. Via the proportion of the two compositions it is possible to increment steplessly the amount of the active compound which is present only in the second composition and, where appropriate, in the at least one further composition. In this way, therefore, it is readily possible to dose any desired amount and any desired proportion of the first to the second active compound and, where appropriate, of the first active compound to a further active compound, without altering the concentration of the first active compound.

The first active compound may be at least one insulin. The second active compound may be at least one GLP-1 agonist. Preference is given to a medicament which comprises a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, the first pharmaceutical composition comprising at least one insulin, and the second pharmaceutical composition comprising at least one insulin and at least one GLP-1 agonist, and the at least one further pharmaceutical composition comprising at least one insulin and at least one further active compound.

The first composition preferably comprises as active compound only the at least one insulin.

The further active compound may be any desired active compound. More particularly the further active compound is an active compound which is used for treating patients with diabetes mellitus (type 1 and/or type 2), including active compounds for treating concomitant disorders of diabetes as well.

The first, second, and, where used, at least one further composition may comprise the insulin in a substantially equal weight fraction or in different weight fractions relative to the total weight of the composition.

It is preferred for the first pharmaceutical composition, the second pharmaceutical composition, and, where used, the at least one further pharmaceutical composition to comprise the insulin in substantially equal weight fractions relative to the total weight of the composition. By this means it is possible to ensure that any desired proportion of the first and second composition and, where appropriate, any desired proportion of the first and at least one further compositions can be used, the dosing of the insulin taking place via the total amount of the compositions administered. Via the proportion of the two compositions it is possible to increment steplessly the amount of the active compound which is present only in the second composition and, where appropriate, in the at least one further composition. In this way, therefore, it is readily possible to dose any desired amount and any desired proportion of insulin to GLP-1 agonist and, where appropriate, of insulin to a further active compound, without altering the concentration of the at least one insulin.

In the present invention, "substantially equal weight fractions" of an active compound in two compositions means that one of the two compositions contains the active compound in a weight fraction which is, for example, not more than 10%, not more than 5%, not more than 1% or not more than 0.1% higher than its weight fraction in the other composition.

The first active compound may also be at least one GLP-1 agonist. The second active compound may be at least one insulin. Preference is given to a medicament which comprises a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, the first pharmaceutical composition comprising at least one GLP-1 agonist, and the second pharmaceutical composition comprising at least one GLP-1 agonist and at least one insulin, and the at least one

12

further pharmaceutical composition comprising at least one GLP-1 agonist and at least one further active compound.

The first composition preferably comprises as active compound only the at least one GLP-1 agonist.

The first, second, and, where used, at least one further compositions may comprise the GLP-1 agonist in a substantially equal weight fraction or in different weight fractions relative to the total weight of the composition. It is preferred for the first pharmaceutical composition, the second pharmaceutical composition and, where used, the at least one further pharmaceutical composition to comprise the at least one GLP-1 agonist in substantially equal weight fractions relative to the total weight of the composition.

Accordingly the present invention provides a medicament which exhibits a number of advantages over compositions of the prior art comprising separate compositions each containing one active compound, more particularly an insulin or a GLP-1 agonist, said advantages including the following:

the ratio of the first active compound to the second active compound and, where appropriate, of the first active compound to the at least one further active compound can be chosen freely by the user.

Since the first active compound is present in all of the compositions, more particularly in equal weight fractions, this active compound is not diluted when the first composition is mixed with the second and, where appropriate, further compositions. This is important for active compounds such as insulin, for example, where the pharmacokinetics is influenced by the concentration/dilution.

The injection volume is reduced (see example 12). Hence there is a reduction in the dilution of the second active compound (e.g., a GLP-1 agonist) and, where appropriate, of a further active compound.

The invention additionally provides a kit comprising a medicament of the invention. The kit of the invention may be intended for use by medical staff or by persons without specialist medical training, more particularly by the patients themselves or helpers such as relatives. In the kit of the invention, the individual pharmaceutical compositions comprising the medicament of the invention are assembled in separate packs, and so the patient is able to select the composition appropriate to the current requirement and to administer an amount in line with that requirement. The kit of the invention comprises, for example, the medicament of the invention in the form of a set of syringes, glass ampoules and/or pens which comprise a composition of the invention.

There are a variety of ways in which the medicament of the invention can be administered. The medicament may be administered parenterally. The medicament may be injected, with the possible use of injection systems with or without needles. Furthermore, the medicament may be administered by inhalation. In this case it is possible for liquid compositions to be inhaled, or the compositions can be inhaled in the form of powder. Furthermore, the medicament of the invention may be administered as a spray, more particularly as a nasal spray. In addition the medicament of the invention may be administered by a transdermal system. The skilled worker is aware of these methods of administration and is able to formulate the medicament of the invention in such a way that it can be effectively administered by one of these methods of administration. The compositions of the medicament of the invention are preferably liquid. It is preferred, furthermore, for the medicament of the invention to be administered parenterally, more particularly by injection.

The present invention further provides a device for administering the medicament of the invention. This device

US 9,526,764 B2

13

comprises the pharmaceutical compositions which are encompassed by the medicament of the invention, in separate containers, and allows the pharmaceutical compositions to be dosed independently of one another. The device of the invention may be a device for parenteral administration. The device of the invention may be a device for injection with or without needles. Furthermore, the device may be a device for inhalation, in which case liquid compositions are inhaled, or the compositions can be inhaled in the form of powder. Moreover, the device may be a device for administering a spray, more particularly a nasal spray. In addition, the device may be a transdermal administration system. It is preferred for the device of the invention to be a device for parenteral administration, more particularly an injection device.

“Compounding” is a term which is known to the skilled worker and which in pharmacology identifies the finishing treatment, such as portioning and packing, for example, of medicaments for use by the end user. In the present specification, “compounded” or “compounding” means more particularly that the pharmaceutical compositions of the invention are packaged in a suitable way in a therapeutically effective amount to allow the herein-described selection of at least one of the compositions of the medicament of the invention for the desired dosing of the at least one insulin and of the at least one GLP-1 agonist. Intended more particularly is a parenteral administration, preferably an injection, more preferably subcutaneous injection. Suitable packaging is, for example, a syringe or a glass vessel with a suitable closure, from which, as required, individual therapeutically active doses can be taken. Likewise suitable are injection pens for the administration of insulin, comprising a container (e.g., a cartridge) which contains a pharmaceutical composition of the invention.

“Formulating” or “formulation” is a term which is known to the skilled worker and which, in the field of pharmacology, refers to the production of medicaments and medicament compositions, and their preparation with excipients. In the present specification “formulating” or “formulation” means more particularly that the composition of the invention is provided in a suitable form which allows administration of a therapeutically effective amount of the active compounds. More particularly a formulation is intended for parenteral administration, preferably for injection, more preferably for subcutaneous injection.

In the present invention the term “GLP-1 agonist” include GLP-1, analogs and derivatives thereof, exendin-3 and analogs and derivatives thereof, and exendin-4 and analogs and derivatives thereof. The compositions of the invention comprise one or more selected independently of one another from the group consisting of glucagon-like peptide-1 (GLP-1), analogs and derivatives of GLP-1, exendin-3, analogs and derivatives of exendin-3, exendin-4, analogs and derivatives of exendin-4, and pharmacologically tolerable salts thereof. Also included are substances which exhibit the biological activity of GLP-1.

GLP-1 analogs and derivatives are described in WO 98/08871, for example; exendin-3, analogs and derivatives of exendin-3, and exendin-4 and analogs and derivatives of exendin-4 can be found in WO 01/04156, WO 98/30231, U.S. Pat. No. 5,424,286, in EP application 99 610043.4, in WO 2004/005342 and WO 04/035623. These documents are included herein by reference. The exendin-3 and exendin-4 described in these documents, and the analogs and derivatives thereof that are described there, can be used in the compositions of the present invention as GLP-1 agonists. It is also possible to use any desired combinations of the

14

exendin-3 and exendin-4 described in these documents, and the analogs and derivatives described therein, as GLP-1 agonists. The at least one GLP-1 agonist is preferably independently selected from the group consisting of exendin-4, analogs and derivatives of exendin-4, and pharmacologically tolerable salts thereof.

A further preferred GLP-1 agonist is an analog of exendin-4 selected from a group consisting of:

H-desPro³⁶-exendin-4-Lys₆-NH₂,
H-des(Pro^{36,37})-exendin-4-Lys₆-NH₂,
H-des(Pro^{36,37})-exendin-4-Lys₅-NH₂, and pharmacologically tolerable salts thereof.

A further preferred GLP-1 agonist is an analog of exendin-4 selected from a group consisting of:

desPro³⁶[Asp²⁸]exendin-4 (1-39),
desPro³⁶[IsoAsp²⁸]exendin-4 (1-39),
desPro³⁶[Met(O)¹⁴, Asp²⁸]exendin-4 (1-39),
desPro³⁶[Met(O)¹⁴, IsoAsp²⁶]exendin-4 (1-39),
desPro³⁶[Trp(O₂)²⁵, Asp²⁸]exendin-2 (1-39),
desPro³⁶[Trp(O₂)²⁶, IsoAsp²⁸]exendin-2 (1-39),
desPro³⁶[Met(O)¹⁴Trp(O₂)²⁵, Asp²⁸]exendin-4 (1-39),
desPro³⁶[Met(O)¹⁴Trp(O₂)²⁶, IsoAsp²⁸]exendin-4 (1-39),
and pharmacologically tolerable salts thereof.

A further preferred GLP-1 agonist is an analog of exendin-4 selected from a group as described in the paragraph above in which the peptide-Lys₆-NH₂ has been attached at the C-termini of the analogs of exendin-4.

A further preferred GLP-1 agonist is an analog of exendin-4 selected from a group consisting of:

H-(Lys)₆-des Pro³⁶[Asp²⁶]exendin-4(1-39)-Lys₆-NH₂
des Asp²⁸Pro³⁶, Pro³⁷, Pro³⁸ exendin-4(1-39)-NH₂,
H-(Lys)₆-des Pro³⁶, Pro³⁷, Pro³⁸[Asp²⁶]exendin-4(1-39)-NH₂,
H-Asn-(Glu)₅ des Pro³⁶, Pro³⁷, Pro³⁸[Asp²⁸]exendin-4(1-39)-NH₂,
des Pro³⁶, Pro³⁷, Pro³⁸[Asp²⁸]exendin-4(1-39)-(Lys)₆-NH₂,
H-(Lys)₆-des Pro³⁶, Pro³⁷, Pro³⁸[Asp²⁸]exendin-4(1-39)-(Lys)₆-NH₂,
H-Asn-(Glu)₅-des Pro³⁶, Pro³⁷, Pro³⁸[Asp²⁸]exendin-4(1-39)-(Lys)₆-NH₂,
H-(Lys)₈-des Pro³⁶[Trp(O₂)²⁵, Asp²⁸]exendin-4(1-39)-Lys₆-NH₂,
H-des Asp²⁸ Pro³⁶, Pro³⁷, Pro³⁸[trp(O₂)²⁶]exendin-4(1-39)-NH₂,
H-(Lys)₆-des Pro³⁶, Pro³⁷, Pro³⁸[Trp(O₂)²⁵, Asp²⁸]exendin-4(1-39)-NH₂,
H-Asn-(Glu)₅-des Pro³⁶, Pro³⁷, Pro³⁸[Trp(O₂)²⁶, Asp²⁸]exendin-4(1-39)-NH₂,
des Pro³⁶, Pro³⁷, Pro³⁸[trp(O₂)²⁶, Asp²⁸]exendin-4(1-39)-(Lys)₆-NH₂,
H-(Lys)₆-des Pro³⁶, Pro³⁷, Pro³⁸[Trp(O₂)²⁵, Asp²⁸]exendin-4(1-39)-(Lys)₆-NH₂,
H-Asn-(Glu)₅-des Pro³⁶, Pro³⁷, Pro³⁸[Trp(O₂)²⁶, Asp²⁸]exendin-4(1-39)-(Lys)₆-NH₂,
H-(Lys)₈-des Pro³⁶[Met(O)¹⁴, Asp²⁸]exendin-4(1-39)-Lys₈-NH₂,
des Met(O)¹⁴ Asp²⁸ Pro³⁶, Pro³⁷, Pro³⁸ exendin-4(1-39)-NH₂,
H-(Lys)₆-des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Asp²⁸]exendin-4(1-39)-NH₂,
H-Asn-(Glu)₅-des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Asp²⁸]exendin-4(1-39)-NH₂,
des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Asp²⁸]exendin-4(1-39)-(Lys)₆-NH₂,
H-(Lys)₆-des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Asp²⁸]exendin-4(1-39)-Lys₆-NH₂,

US 9,526,764 B2

15

H-Asn-(Glu)₅-des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Asp²⁸]ex-
 endin-4(1-39)-(Lys)₆-NH₂,
 H-(Lys)₆-des Pro³⁶[Met(O)¹⁴, Trp(O₂)²⁶, Asp²⁸]exendin-4
 (1-39)-Lys₆-NH₂,
 des Asp²⁸ Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Trp(O₂)²⁸]exendin-
 4(1-39)-NH₂,
 H-(Lys)₆-des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Trp(O₂)²⁶,
 Asp²⁸]exendin-4(1-39)-NH₂,
 H-Asn-(Glu)₅-des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Asp²⁸]ex-
 endin-4(1-39)-NH₂,
 des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Trp(O₂)²⁶, Asp²⁸]exen-
 din-4(1-39)-(Lys)₆-NH₂,
 H-(Lys)₆-des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Trp(O₂)²⁵,
 Asp²⁸]exendin-4(1-39)-(Lys)₆-NH₂,
 H-Asn-(Glu)₅-des Pro³⁶, Pro³⁷, Pro³⁸[Met(O)¹⁴, Trp(O₂)²⁸,
 Asp²⁸]exendin-4(1-39)-(Lys)₆-NH₂, and pharmacologically
 tolerable salts thereof.

A further preferred GLP-1 agonist is selected from a
 group consisting of Arg³⁴, Lys²⁶(N^ε(γ-glutamyl(N^α-hexade-
 canoyl)))GLP-1(7-37)[liraglutide] and a pharmacologically
 tolerable salt thereof.

A further preferred GLP-1 agonist is AVE0010. AVE0010
 has the sequence of desPro³⁶ exendin-4(1-39)-Lys₆-NH₂.
 This substance is published as SEQ ID No: 93 in WO
 01/04156. Preference is also given to pharmacologically
 tolerable salts of AVE0010.

The term “at least one GLP-1 agonist” includes combi-
 nations of the herein-described GLP-1 agonists which are
 used in the compositions of the invention, examples being
 any desired combinations of two or more GLP-1 agonists
 selected from the GLP-1 agonists described herein.

The at least one GLP-1 agonist is further preferably
 independently selected from exendin-4, Pro³⁶exendin-4(1-
 39)-Lys₆-NH₂, and Arg³⁴, Lys²⁶(N^ε(γ-glutamyl(N^α-hexade-
 canoyl)))GLP-1(7-37)[liraglutide], and pharmacologically
 tolerable thereof.

The compositions of the invention contain the GLP-1
 agonist in an amount of 10 μg/ml to 20 mg/ml, preferably 25
 μg/ml to 15 mg/ml. For the acidic to neutrally dissolved
 GLP-1 agonists the figures are preferably 20 μg/ml to 300
 μg/ml, and for the neutral to basic agonists they are prefer-
 ably 500 μg/ml to 10 mg/ml. For exendin-4 analogs, 20
 μg/ml to 150 μg/ml are preferred.

In the present specification the term “insulin” encom-
 passes not only unmodified insulins but also insulin analogs,
 insulin derivatives, and insulin metabolites. The composi-
 tions of the invention comprise one or more independently
 selected from the group consisting of insulins (e.g., unmodi-
 fied insulins), insulin analogs, insulin derivatives, and insu-
 lin metabolites, and any desired combinations thereof.

The at least one insulin may independently be selected
 from the group consisting of bovine insulins, analogs,
 derivatives, and metabolites thereof, porcine insulins, ana-
 logs, derivatives, and metabolites thereof, and human insu-
 lins, analogs, derivatives, and metabolites thereof. Prefer-
 ably the at least one insulin is independently selected from
 human insulins, analogs, derivatives, and metabolites
 thereof.

Furthermore, an insulin of the invention may be selected
 independently from unmodified insulins, more particularly
 from bovine insulins, porcine insulins, and human insulins.

The at least one insulin may independently be selected
 from the group consisting of bovine insulins, porcine insu-
 lins, and human insulins. More preferably the at least one
 insulin is independently selected from human insulins. An

16

insulin of the invention may be selected from unmodified
 insulins, more particularly from bovine insulins, porcine
 insulins, and human insulins.

Insulin derivatives of the invention are derivatives of a
 naturally occurring insulin and/or an insulin analog, which
 are obtained by chemical modification. The chemical modi-
 fication may consist, for example, in the addition of one or
 more defined chemical groups onto one or more amino
 acids.

Insulin analogs which are described in EP 0 214 826, EP
 0 375 437, EP 0 678 522, EP 0 885 961, EP 0 419 504, WO
 92/00321, German patent applications 10 2008 003 568.8
 and 10 2003 003 566.1, and EP-A 0 368 187 may be part of
 the compositions of the invention. The documents EP 0 214
 826, EP 0 375 437, EP 0 678 522, EP 0 419 504, WO
 92/00321, and EP-A 0 368 187 are included herein by
 reference.

One preferred insulin analog of the invention may be
 selected from the group consisting of Gly(A21)-Arg(B31)-
 Arg(B32) human insulin (insulin glargine, Lantus); Arg
 (A0)-His(A8)-Glu(A15)-Asp(A18)-Gly(A21)-Arg(B31)-
 Arg(B32) human insulin amide, Lys(B3)-Glu(B29) human
 (insulin; Lys^{B28}Pro^{B29} human insulin (insulin lyspro), B28
 Asp human insulin (insulin aspart), human insulin in which
 proline in position B28 has been substituted by Asp, Lys,
 Leu, Val or Ala and where Lys in position B29 may be
 substituted by Pro; AlaB26 human insulin; des(B28-B30)
 human insulin; des(B27) human insulin or B29Lys(ε-tetra-
 decanoyl), des(B30) human insulin (insulin detemir).

A preferred insulin derivative of the invention may be
 selected from the group consisting of B29-N-myristoyl-des
 (B30) human insulin, B29-N-palmitoyl-des(B30) human
 insulin, B29-N-myristoyl human insulin, B29-N-palmitoyl
 human insulin, B28-N-myristoyl Lys^{B28}Pro^{B29} human insu-
 lin, B28-N-palmitoyl-Lys^{B28}Pro^{B29} human insulin, B30-N-
 myristoyl-Thr^{B29}Lys^{B30} human insulin, B30-N-palmitoyl-
 Thr^{B29}Lys^{B30} human insulin. B29-N-(N-palmitoyl-Y-
 glutamyl)-des(B30) human insulin, B29-N-(N-lithocholyl-
 Y-glutamyl)-des(B30) human insulin, B29-N-(ω-
 carboxyheptadecanoyl)-des(B30) human insulin, and B29-
 N-(ω-carboxyheptadecanoyl) human insulin.

A more highly preferred insulin derivative of the inven-
 tion is selected from the group consisting of Gly(A21)-Arg
 (B31)-Arg(B32) human insulin, Ly^{B28}Pro^{B29} human insulin
 (insulin lyspro), B28 Asp human insulin (insulin aspart),
 B29Lys(ε-tetradecanoyl), desB30 human insulin (insulin
 detemir).

The term “at least one insulin” includes combinations of
 the herein-described insulins, analogs, derivatives, and
 metabolites thereof which are used in the compositions of
 the invention, e.g., any desired combinations of two or more
 selected from the herein-described insulins, analogs, deriva-
 tives, and metabolites.

The compositions of the invention contain 60-6000 nmol/
 ml, preferably 240-3000 nmol/ml, of an insulin as defined
 herein. Depending on the insulin used, a concentration of
 240-3000 nmol/ml corresponds approximately to a concen-
 tration of 1.4-35 mg/ml or 40-500 units/ml.

In the 2 to 10, preferably 3 to 5, pens cover all system, the
 compositions are in the range from 20 μg/ml of GLP-1
 agonist and 100 U/ml of insulin to 300 μg/ml of GLP-1
 agonist and 500 U/ml of insulin. Preference is given to the
 following concentration ranges: 25 μg/ml and 100 U/ml, 33
 μg/ml and 100 U/ml, 40 μg/ml and 100 U/ml, 66 μg/ml and
 100 U/ml, and 75 μg/ml and 100 U/ml.

The desired dosage range of the insulin is in particular a
 dosage with a synergistic effect. Here the values are 5 to 100

US 9,526,764 B2

17

U, preferably 15 to 80 U. For the GLP-1 agonist the values for the dosage range are 5 µg to 2 mg, preferably 10 µg to 1.8 mg, more preferably 10 µg to 30 µg.

[More Precise Details Here on the Amounts Used and on the Dosages]

The preferred presentation form of the pharmaceutical compositions of the present invention is that of liquid compositions suitable in particular for parenteral administration, more preferably for injection, most preferably for subcutaneous injection. In particular the pharmaceutical composition of the present invention is suitable for injection once daily.

The pharmaceutical composition of the present invention may have an acidic or physiological pH. An acidic pH range is situated preferably in the range of pH 1-6.8, more preferably pH 3.5-6.8, even more preferably pH 3.5-4.5, most preferably at a pH of about 4.0-4.5. A physiological pH is situated preferably in the range of pH 4.0-8.5, more preferably pH 5.0 to 8.5, even more preferably pH 6.0 to 8.5.

The composition of the invention may comprise a suitable preservative. Examples of suitable preservatives include phenol, m-cresol, benzyl alcohol and/or p' esters.

The composition of the invention may further comprise a suitable buffer. Buffer substances which can be used, particularly for setting a pH level between about 4.0 and 8.5, include, for example, sodium acetate, sodium citrate, sodium phosphate, etc. Otherwise, physiologically unobjectionable dilute acids (typically HCl) or alkalis (typically NaOH) are suitable for setting the pH level. Preferred concentrations of the buffers and also of corresponding salts are in the range of 5-250 mM, more preferably in the range of 10-100 mM.

The composition of the invention may comprise zinc ions. The concentration of the zinc ions is preferably in the range from 0 µg/ml to 500 µg/ml, more preferably from 5 µg to 200 µg of zinc/ml.

The composition of the invention may further comprise suitable isotonicity agents. Suitable examples include glycerol, dextrose, lactose, sorbitol, mannitol, glucose, NaCl, calcium compounds or magnesium compounds such as CaCl₂, etc. Glycerol, dextrose, lactose, sorbitol, mannitol, and glucose are typically in the range of 100-250 mM, NaCl in a concentration of up to 150 mM.

The composition of the invention may further comprise a surfactant. A surfactant may greatly increase the stability of acidic insulin compositions. Using surfactant it is even possible to prepare compositions which guarantee the superior stability with respect to hydrophobic aggregation nuclei over a number of months with temperature exposure.

The surfactant is preferably selected from the group consisting of partial and fatty acid esters and ethers of polyhydric alcohols such as of glycerol and of sorbitol, and polyols, the partial and fatty acid esters and ethers of glycerol and of sorbitol being selected from a group containing Span®, Tween®, Myrj®, Brij®, and Cremophor®; and the polyols being selected from the group of polypropylene glycols, polyethylene glycols, poloxamers, polysorbates, Pluronic, and Tetronics. Preferred concentrations of the surfactants are in the range of 5-200 µg/ml, preferably of 5-120 µg/ml and more preferably of 20-75 µg/ml.

The composition of the invention may further comprise other additives such as, for example, salts, which retard the release of at least one insulin.

One particularly preferred subject of the invention is a medicament as described herein comprising at least one insulin independently selected from Lys^{B28}Pro^{B29} human insulin (insulin lyspro), B28 Asp human insulin (insulin aspart), B29Lys(ε-tetradecanoyl), desB30 human insulin

18

(insulin detemir), and insulin glargine (Gly(A21)-Arg(B31)-Arg(B32) human insulin), and comprising AVE0010 and/or a pharmacologically tolerable salt thereof. A further particularly preferred subject is a medicament as described herein comprising insulin glargine (Gly(A21)-Arg(B31)-Arg(B32) human insulin) and AVE0010 (des Pro³⁶exendin-4(1-39)-Lys₆-NH₂) and/or a pharmacologically tolerable salt thereof. The compositions of these particularly preferred medicaments preferably have an acidic pH of 1-6.8, more preferably pH 3.5-6.8, even more preferably pH 3.5-5.0, most preferably a pH of about 4.0 to 4.5. In addition, the compositions of these particularly preferred medicaments may comprise a surfactant as described herein.

A further subject of the invention is a combination of insulin glargine (Gly(A21)-Arg(B31)-Arg(B32) human insulin) and AVE0010 (des Pro³⁶exendin-4(1-39)-Lys₆-NH₂) and/or a pharmacologically tolerable salt thereof.

The invention further provides a method of treating a patient with a kit or medicament of the invention as described herein.

The method of the invention for treating a patient comprises the administration of a medicament of the invention comprising at least one insulin and at least one GLP-1 agonist, the medicament being formulated and/or compounded such that it contains the insulin and the GLP-1 agonist each in a predetermined amount and can be administered in a dose adapted to the individual requirement of a patient.

More particularly the method comprises the administration of a medicament comprising a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, each comprising at least one insulin and at least one GLP-1 agonist, and comprising the at least one insulin and/or the at least one GLP-1 agonist in different weight fractions relative to the total weight of the composition, said method comprising:

- (a) selecting a dose of the at least one insulin that is to be administered,
- (b) selecting a dose of the at least one GLP-1 agonist that is to be administered,
- (c) selecting a composition, from the first, second, and, where used, at least one further compositions of the medicament that comprises the doses from (a) and (b) in a concentration such that the doses from (a) and (b) are present in the same volume, and
- (d) determining and administering an amount which corresponds to the doses from (a) and (b).

The dose according to step (a) and/or step (b) is determined according to the individual requirement of the patients.

Step (c) of the treatment method of the invention can be carried out on the basis of a table. This table may be part of the medicament of the invention. Example 11 contains an example of a table of the invention.

The method of treating a patient may more particularly comprise the administration of a medicament, the medicament comprising a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, the first pharmaceutical composition comprising at least one first active compound, and the second pharmaceutical composition comprising at least one first active compound and at least one second active compound, the at least one further pharmaceutical composition comprising at least one first active compound and at least one further active compound, and the method comprising the following steps:

US 9,526,764 B2

19

- (i) selecting a dose of the at least one first active compound that is to be administered, and determining the total amount of the first, second, and, where used, at least one further composition, so that the selected dose of the at least one first active compound is present in the total amount,
- (ii) selecting a dose of the at least one second active compound that is to be administered and determining the amount of the second composition so that the selected dose of the at least one second active compound is present in the amount of the second composition,
- (iii) where appropriate, selecting a dose of the at least one further active compound that is to be administered, and determining the amount of the at least one further composition so that the selected dose of the at least one further active compound is present in the amount of the at least one further composition,
- (iv) administering an amount of the first composition to the patient, the administered amount corresponding to the total amount as per step (i) minus the amount of the second composition as per step (ii), and, where appropriate, minus the amount of the at least one further composition as per step (iii), and
- (v) administering the amount of the second composition that was determined in step (ii), and, where appropriate, the amount of the at least one further composition that was determined in step (iii), to the patient.

The first active compound may be an insulin, and the second active compound may be a GLP-1 agonist. Hence the method of treating a patient may comprise more particularly the administration of a medicament, the medicament comprising a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, the first pharmaceutical composition comprising at least one insulin, and the second pharmaceutical composition comprising at least one insulin and at least one GLP-1 agonist, and the at least one further pharmaceutical composition comprising at least one insulin and at least one further active compound, and the method comprising the steps of:

- (i) selecting a dose of the at least one insulin that is to be administered, and determining the total amount of the first, second, and, where used, at least one further composition, so that the selected dose of the at least one insulin is present in the total amount,
- (ii) selecting a dose of the at least one GLP-1 agonist that is to be administered and determining the amount of the second composition so that the selected dose of the at least one GLP-1 agonist is present in the amount of the second composition,
- (iii) where appropriate, selecting a dose of the at least one further active compound that is to be administered, and determining the amount of the at least one further composition so that the selected dose of the at least one further active compound is present in the amount of the at least one further composition,
- (iv) administering an amount of the first composition to the patient, the administered amount corresponding to the total amount as per step (i) minus the amount of the second composition as per step (ii), and, where appropriate, minus the amount of the at least one further composition as per step (iii), and
- (v) administering the amount of the second composition that was determined in step (ii), and, where appropriate, the amount of the at least one further composition that was determined in step (iii), to the patient.

20

The first active compound may be a GLP-1 agonist, and the second active compound may be an insulin. Hence the method of treating a patient may comprise more particularly the administration of a medicament, the medicament comprising a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, the first pharmaceutical composition comprising at least one GLP-1 agonist, and the second pharmaceutical composition comprising at least one GLP-1 agonist and at least one insulin, and the at least one further pharmaceutical composition comprising at least one GLP-1 agonist and at least one further active compound, and the method comprising the steps of:

- (i) selecting a dose of the at least one GLP-1 agonist that is to be administered, and determining the total amount of the first, second, and, where used, at least one further composition, so that the selected dose of the at least one GLP-1 agonist is present in the total amount,
- (ii) selecting a dose of the at least one insulin that is to be administered and determining the amount of the second composition so that the selected dose of the at least one insulin is present in the amount of the second composition,
- (iii) where appropriate, selecting a dose of the at least one further active compound that is to be administered, and determining the amount of the at least one further composition so that the selected dose of the at least one further active compound is present in the amount of the at least one further composition,
- (iv) administering an amount of the first composition to the patient, the administered amount corresponding to the total amount as per step (i) minus the amount of the second composition as per step (ii), and, where appropriate, minus the amount of the at least one further composition as per step (iii), and
- (v) administering the amount of the second composition that was determined in step (ii), and, where appropriate, the amount of the at least one further composition that was determined in step (iii), to the patient.

Steps (i), (ii) and/or (iii) may be carried out on the basis of at least one table, which may be part of the medicament. For each of steps (i), (ii), and (iii) independently of one another a table may be provided.

The treatment method of the invention may be used more particularly for treating patients with diabetes, more particularly with diabetes type 1 or II. Preferably the method is used to adjust the fasting, postprandial and/or postabsorptive blood glucose concentration, for improving glucose tolerance, for preventing hypoglycemia, for preventing loss of function of the pancreatic β cells, for weight loss and/or for preventing weight gain.

The invention further provides a method of preparing a medicament of the invention, comprising formulating and/or compounding, such that it contains the insulin and the GLP-1 agonist each in a predetermined amount and can be administered in a dose adapted to the individual requirement of a patient. In the preparation method the medicament is preferably formulated and compounded such that one of the herein-described medicaments of the invention can be obtained, as for example a medicament of the invention comprising a first pharmaceutical composition and a second pharmaceutical composition, and, optionally, at least one further pharmaceutical composition, each comprising at least one insulin and at least one GLP-1 agonist, and comprising the at least one insulin and/or the at least one GLP-1 agonist in different weight fractions relative to the total weight of the composition.

US 9,526,764 B2

21

The invention is illustrated by the following figures and the following example, which do not restrict the invention in any way whatsoever.

KEY TO THE FIGURES

FIG. 1: Study design for oral glucose tolerance test.

FIG. 2: OGTT in the dog: Effect of insulin glargine relative to placebo.

FIG. 3: OGTT in the dog: Effect of AVE0010 relative to placebo.

FIG. 4: OGTT in the dog: Effect of an AVE0010/insulin glargine combination on blood glucose level.

FIG. 5: OGTT in the dog: Effect of an AVE0010/insulin glargine combination on plasma insulin and the c-peptide level.

FIG. 6: OGTT in the dog: Effect of a dose lowering of AVE0010 with different proportions relative to insulin glargine in the combined formulation.

FIG. 7: Effect of an AVE0010/insulin glargine combination on blood glucose in the diabetic db/db mouse.

FIG. 8: Effect of an AVE0010/insulin glargine combination in the oral glucose tolerance test in the diabetic db/db mouse.

FIG. 9: Effect of an AVE0010/insulin glargine combination on cytokine- and lipotoxicity-induced β -cell apoptosis in vitro.

FIG. 10: The "3 pens cover all" system.

EXAMPLES

Example 1

Model: Oral glucose tolerance test (OGTT) in healthy dogs: Comparison of the insulin glargine/AVE0010 combination with the two individual active compounds.

Animals

Male normoglycemic beagles

Bodyweight: ~15 kg

Number per group: n=6

Study Design (See FIG. 1)

Individual subcutaneous injections of placebo or test formulation at time 0

2 oral administrations of glucose, at 2 g of glucose/kg of bodyweight, at times 30 min and 5 h

Blood samples are taken to determine blood glucose, plasma insulin, and c-peptide

Group Division (n=6)

Placebo (Lantus placebo formulation without API)

Insulin glargine (0.3 IU/kg s.c., equivalent to 1.8 nmol/kg). Insulin glargine is Gly(A21)-Arg(B31)-Arg(B32) human insulin.

AVE0010 (10 μ g/kg s.c. in Lantus placebo formulation, equivalent to 2 nmol/kg). AVE0010 is des Pro³⁶exendin-4(1-39)-Lys₆-NH₂.

AVE0010/insulin glargine combination (10 μ g/kg AVE0010/0.3 IU/kg insulin glargine s.c.)

Example 2

OGTT in the Dog: Effect of Insulin Glargine Relative to Placebo

The Experiment was Carried Out in Accordance with the Protocol Described in Example 1.

repeated OGTT (2 g/kg p.o.)

male beagle, n=6

22

mean \pm Sem

placebo=Lantus placebo

insulin glargine (0.3 U/kg s.c.)

Result: The data are shown in FIG. 2. The single administration of insulin glargine does not prevent the OGTT-induced increase in blood glucose. Insulin glargine reinforces the expected delayed lowering of blood glucose concentration in the postabsorptive phase.

Example 3

OGTT in the Dog: Effect of AVE0010 Relative to Placebo

The experiment was carried out in accordance with the protocol described in example 1.

repeated OGTT (2 g/kg p.o.)

male beagle, n=6

mean \pm Sem

placebo=Lantus placebo

AVE0010 (10 μ g/kg s.c.)

Result: The data are shown in FIG. 3. AVE0010 prevents the OGTT-induced postprandial increase in blood glucose almost completely. There is no effect on the glucose concentration in the postabsorptive phase. This example shows that the effect of AVE0010 on the OGTT-induced postprandial increase in blood glucose is complementary to the blood sugar-lowering effect of insulin glargine in the postabsorptive phase.

Example 4

OGTT in the Dog: Effect of an AVE0010/Insulin Glargine Combination on the Blood Glucose Level

The experiment was carried out in accordance with the protocol described in example 1.

repeated OGTT (2 g/kg p.o.)

male beagle, n=6

mean \pm Sem

placebo=Lantus placebo

AVE0010 (10 μ g/kg s.c.)

Insulin glargine (0.3 U/kg s.c.)

AVE+Lan (=premix of 10 μ g/kg of AVE0010 and 0.3 U/kg of insulin glargine in one formulation)

Result: The data are shown in FIG. 4. The combination has the same action on the postprandial glucose increase as AVE0010 (cf. example 3). The hypoglycemic effect of insulin glargine in the postabsorptive phase is likewise present, but attenuated (cf. example 2). This is a synergistic effect of insulin glargine and AVE0010, since AVE0010 alone has no effect on the level of glucose, which has fallen again following administration of glucose, and insulin glargine on its own has no effect on the postprandial glucose level.

Example 5

OGTT in the Dog: Effect of an AVE0010/Insulin Glargine Combination on the Plasma Insulin and the c-Peptide Level

The experiment was carried out in accordance with the protocol described in example 1.

repeated OGTT (2 g/kg p.o.)

a male beagle, n=6

a mean \pm Sem

US 9,526,764 B2

23

placebo=Lantus placebo
 AVE0010 (10 µg/kg s.c.)
 glargine (0.3 U/kg s.c.)
 AVE+Lan (=premix of 10 µg/kg of AVE0010 and 0.3 U/kg
 of insulin glargine in one formulation)

The C-peptide is released in the course of the conversion of proinsulin to insulin, and serves as a marker for the secretion of insulin by the pancreatic β-cells. In a glucose loading test, the c-peptide can be used to determine the response capacity of the pancreas.

Result: The data are shown in FIGS. 5a and 5b. In the combination group, the postprandial reduction in insulin is followed by an increased postabsorptive insulin glargine level. C-peptide levels for the combination correspond to the insulin curve of AVE0010 during the prandial phases, and of insulin glargine during the postabsorptive phase.

Example 6

OGTT in the Dog: Effect of a Dose Lowering of
 AVE0010 with Different Proportions to Insulin
 Glargine in the Combined Formulation

The experiment was carried out in accordance with the protocol described in example 1.

repeated OGTT (2 g/kg p.o.)
 male beagle, n=11/6/6/6
 mean±Sem

control=Lantus placebo

AVE+Lan (=premix of 0.15 to 1.0 µg/kg of AVE0010 and 0.3 U/kg of insulin glargine in one formulation). In examples 2 to 5, AVE0010 concentrations of 10 µg/kg were used.

Result: The data are shown in FIG. 6. A reduction in the AVE0010 dose from 10 µg/kg (cf. in particular example 4) to 1 µg/kg (i.e., by a factor of 10), and the resultant increase in the proportion of insulin glargine to AVE0010, has no effect on the synergistic activity of the combination of AVE0010 with insulin glargine (cf. in particular example 4). Only at significantly smaller AVE0010 doses does the effect of the combination approach the effect of insulin glargine alone (of, in particular FIG. 2). The AVE0010 dose may therefore be varied at least within one order of magnitude (i.e., by a factor of at least 10) without loss of the synergistic effect.

Example 7

Model: Diabetic, Insulin-Resistant db/db Mouse: Comparison of the Insulin Glargine/AVE0010 Combination with the Two Individual Active Compounds.

Animals

Female db/db mouse
 Age: 10-11 weeks
 Number per group: n=10

Study Design

Individual subcutaneous injection of placebo or test formulation

Taking of blood samples to determine blood glucose

Group Division

Placebo (=Lantus placebo formulation without API)
 AVE0010 (10 µg/kg s.c.)
 Insulin glargine (5 IU/kg s.c.)
 AVE0010/insulin glargine combination (premix of 10 µg/kg of AVE0010 plus 5 IU/kg of insulin glargine s.c.)

24

Example 8

Effect of an AVE0010/Insulin Glargine
 Combination on Blood Glucose in the Diabetic
 db/db Mouse

The experiment was carried out in accordance with the protocol described in example 7.

Female db/db mouse, 10 weeks

n=10, mean±Sem

Vehicle=Lantus placebo

AVE0010 (10 µg/kg sc)

Lentils (5 U/kg sc)

AVE0010/insulin glargine (=premix of AVE0010 10 µg/kg and insulin glargine 5 U/kg in one formulation)

Result: The data are shown in FIG. 7. In diabetic db/db mice, the AVE0010/insulin glargine combination produced a more rapid and more pronounced decrease in the blood glucose concentration as compared with the two individual active compounds. Consequently the combination takes diabetic db/db mice closer to normoglycemia than either of the two active compounds alone.

Example 9

Effect of an AVE0010/Insulin Glargine
 Combination in the Oral Glucose Tolerance Test in
 the Diabetic Db/Db Mouse

The experiment was carried out in accordance with the protocol described in example 7. Additionally an OGTT (2 g/kg p.o. @30 min) was carried out

Female db/db mouse, 11 weeks

n=10, mean±Sem

Control=Lantus placebo

AVE0010 (10 µg/kg sc)

Insulin glargine (5 U/kg sc)

AVE0010/insulin glargine (=premix of AVE0010 10 µg/kg and insulin glargine 5 U/kg in one formulation)

Result: The data are shown in FIG. 8. The AVE0010/insulin glargine combination leads to significantly improved glucose tolerance and lower postabsorptive glucose levels.

Example 10

Effect of the AVE0010/Insulin Glargine
 Combination on Cytokine- and
 Lipotoxicity-Induced β-Cell Apoptosis In Vitro

Insulinoma cell line INS-1, rat

Incubation with the test compound for 5 h

Further incubation with a cytokine mix for 22 h (1 ng/mL IFN-γ-4-4 ng/mL IL-1β) or

Further incubation with 0.5 mM FFA for 18 h (palmitates: BSA 31)

The measures used for the apoptosis are the caspase-3 activity and the fragmentation of the cell nuclei, which correlate with apoptosis.

Result: The data are shown in FIG. 9. AVE0010 or insulin glargine (glargine, Glar) alone prevent the apoptosis by ~40-50%. The AVE0010 and insulin glargine combination prevents apoptosis significantly better. On the basis of this synergistic effect, the combination brings about increased protection against cytokine- and lipotoxicity-induced apoptosis.

US 9,526,764 B2

25

Example 11

The “3 pens cover all” system (FIG. 10)

3 premix pens with 3 different predetermined proportions:

Mix A: 100 U of insulin glargine+66.66 µg of AVE0010 per mL

Mix B: 100 U of insulin glargine+40 µg of AVE0010 per mL

Mix C: 100 U of insulin glargine+25 µg of AVE0010 per mL

Use of the 3 premix pens: The table in FIG. 10, representing an example, starts from a therapeutic range of 15 to 80 Upper dose of insulin glargine and 10 to 20 µg of AVE0010. For a particular patient, a dose of insulin glargine to be administered is specified or predetermined. The predetermined dose is looked up in the left-hand column. Where the columns MIX A-MIX C specify a corresponding AVE0010 dose in the range between 10 and 20 µg, the corresponding MIX is selected, dosed, and administered. The ranges are overlapping: for example, in the case of a requirement of 26 to 30 U of insulin glargine, it will be possible to choose MIX A or MIX B (with a higher dose of AVE0010). The same applies to MIX B and C. If, for example, a dose of 50 U of insulin is intended, then 0.5 ml of MIX B or MIX C can be dosed. This dose contains 20 µg (MIX B) or 12.5 µg (MIX C) of AVE0010,

Conclusion: On the assumption that a probable AVE0010 effect is obtained at between 10 and 15 µg, and a therapeutic effect between 15 and 22 µg, almost all patients who take insulin glargine doses of 15-80 U can also obtain therapeutic doses of AVE0010 of between 10 and 20 µg if they use one of the three premix pens which contain three different insulin glargine:AVE0010 ratios (Mix A, B or C). On the basis of the broad range of possible proportions of insulin glargine to AVE0010 (cf. example 6) with a synergistic effect, the proportions in the pens can be tailored such that for each dose of insulin glargine there is a synergistic dose of AVE0010 in at least one pen.

Example 12

This example shows how a combination of two or more active compounds can be formulated in such a way that, when two or more compositions are combined, both active compounds can be administered in any desired amounts and in any desired proportions to one another. It is taken into account here that at least one of the active compounds must not be diluted as a result of the combining (e.g., through mixing directly prior to administration).

In this example, the designations “active A” and “active B” stand for any desired active compounds. In particular, active A is an insulin and active B is a GLP-1 agonist. Active A can also be a GLP-1 agonist, and active B can also be an insulin.

1. Comparative Example

For a combination therapy with an active A (e.g., an insulin) and an active B (e.g., a GLP-1 agonist), a container 1 with a composition with active A at a concentration of a mg/ml, and a container 2 with a composition with active B at a concentration of b mg/ml, are provided.

For the administration of a combination of the two active, a volume V_1 ml from container 1 and a volume V_2 ml from container 2 are mixed.

26

For the dosing of the two actives, at given concentrations a and b, the volumes V_1 and V_2 to be administered are selected in dependence on the amount of the actives A and B to be administered. The volumes V_1 and V_2 of the two actives are determined on the basis of the amount of active, as follows:

Amount of active A: $V_1 \cdot a$ mg

Amount of active B: $V_2 \cdot b$ mg

The concentrations of the actives A and B in the mixture of the two compositions are determined as follows.

Active A: x mg/mL = $V_1 \cdot a / (V_1 + V_2)$

Active B: y mg/mL = $V_2 \cdot b / (V_1 + V_2)$

$V_1 + V_2$ is the total administered volume. This means that the two actives dilute one another. With this system, therefore, it is not possible to keep, for example, the concentration of the active A (e.g., of the insulin) at a predetermined level in the case of varying amounts of active B.

2. Inventive Example

In this example, for a combination therapy with an active A (e.g., an insulin) and an active B (e.g., a GLP-1 agonist), a container 1 with a composition with active A at a concentration of a mg/ml, and a container 2 with a composition with active A at a concentration of a mg/ml and with active B at a concentration of b mg/ml, are provided. The concentration of the active A is therefore the same in both compositions.

For the administration of a combination of the two actives, a volume V_3 ml from container 1 and a volume V_2 ml from container 2 are mixed.

For the dosing of the two actives, at given concentrations a and b, the volumes V_3 and V_2 to be administered are selected in dependence on the amount of the actives A and B to be administered. The volumes V_3 and V_2 of the two actives are determined on the basis of the amount of active, as follows:

Amount of active A: $(V_3 \cdot a + V_2 \cdot a)$ mg

Amount of active B: $V_2 \cdot b$ mg

The concentrations of the actives A and B are determined as follows.

Active A: a mg/mL = $(V_3 \cdot a + V_2 \cdot a) / (V_3 + V_2)$

Active B: z mg/mL = $V_2 \cdot b / (V_3 + V_2)$

$V_3 + V_2$ is the total administered volume. From the above calculation it is evident that the concentration of the active A is always a mg/ml, i.e., is constant, irrespective of what volume ratio V_3/V_2 is being dosed.

Comparing the comparative example (see section 1) with the present inventive example, it is apparent that, for an equal dosing quantity of actives A and B, the total volume required in the inventive example is lower.

For a given dose (amount of active compound) of the active A, the figure in the

comparative example is: $V_1 \cdot a$ mg

In the inventive example it is: $(V_3 \cdot a + V_2 \cdot a)$ mg

US 9,526,764 B2

27

Since the amount of active compound is to be the same in both cases,

$$(V_3 \cdot a + V_2 \cdot a) = V_1 \cdot a$$

$$(V_3 + V_2) \cdot a = V_1 \cdot a$$

$$\text{and } V_3 + V_2 = V_1$$

$$\text{or } V_3 = V_1 - V_2$$

Here, the volume V_2 in which the active B is administered is the same in both cases.

The total volume in the comparative example is
 $V_1 + V_2$

The total volume in the inventive example is $V_3 + V_2$

According to the above equation, for the inventive example it is the case that:

$$V_3 + V_2 = V_1 - V_2 + V_2 = V_1$$

This volume V_1 is smaller than the volume $V_1 + V_2$ of the comparative example.

As a result of the mixing of the composition with actives A and B with the composition with active A, active B is diluted. This dilution is less than the dilution of the active B in the comparative example (i.e., the concentration $b > \text{concentration } z > \text{concentration } y$):

$$b > z$$

$$b > V_2 \cdot b / (V_3 + V_2)$$

$$b > b \cdot V_2 / (V_3 + V_2), \text{ where } V_2 / (V_3 + V_2) \text{ is } < 1, \text{ and}$$

$$z > y$$

$$V_2 \cdot b / (V_3 + V_2) > V_2 \cdot b / (V_1 + V_2)$$

$$1 / (V_3 + V_2) > 1 / (V_1 + V_2)$$

$$1 / (V_1 - V_2 + V_2) > 1 / (V_1 + V_2)$$

$$1 / V_1 > 1 / (V_1 + V_2)$$

Hence the dosing system of the invention for administering variable doses of the actives A (e.g., an insulin) and B (e.g., a GLP-1 agonist) has three advantages over the comparative system:

The concentration of active A (e.g., an insulin) can be kept constant at a predetermined level

Where the doses of actives A and B to be administered are the same, the total volume to be administered is smaller.

The dilution of active B (e.g., the GLP-1 agonist) is less than in the comparative experiment. Accordingly the concentration of active B can be held more easily within a predetermined range.

The present example can be readily extended to medications with three or more active compounds, the first active compound being present in all of the compositions (preferably in equal weight fractions) and there being at least one further active compound in each further composition. The first composition can be mixed with each further composition in the same proportion without the concentration of the active compound in the first composition becoming diluted.

The invention claimed is:

1. A pharmaceutical formulation comprising a combination of:

(a) insulin glargine or a pharmaceutically acceptable salt thereof, and

28

(b) desPro³⁶exendin-4(1-39)-Lys₆-NH₂ (AVE0010) or a pharmaceutically acceptable salt thereof

wherein the concentration of insulin glargine is about 100 units/mL, and the concentration of AVE0010 is about 20 µg/ml to about 150 µg/ml.

2. The pharmaceutical formulation of claim 1, wherein the concentration of insulin glargine is about 100 units/mL, and the concentration of AVE0010 is about 25, 33, 40, 66, or 75 µg/mL.

3. The pharmaceutical formulation of claim 2, wherein the concentration of insulin glargine is about 100 units/mL, and the concentration of AVE0010 is about 25 µg/mL.

4. The pharmaceutical formulation of claim 2, wherein the concentration of insulin glargine is about 100 units/mL, and the concentration of AVE0010 is about 33 µg/mL.

5. The pharmaceutical formulation of claim 2, wherein the concentration of insulin glargine is about 100 units/mL, and the concentration of AVE0010 is about 40 µg/mL.

6. The pharmaceutical formulation of claim 2, wherein the concentration of insulin glargine is about 100 units/mL, and the concentration of AVE0010 is about 66 µg/mL.

7. The pharmaceutical formulation of claim 2, wherein the concentration of insulin glargine is about 100 units/mL, and the concentration of AVE0010 is about 75 µg/mL.

8. A pharmaceutical formulation comprising a combination of:

(a) insulin glargine or a pharmaceutically acceptable salt thereof, and

(b) AVE0010 or a pharmaceutically acceptable salt thereof;

wherein the concentration of insulin glargine is about 100 units/mL, and

wherein insulin glargine and AVE0010 are present in a ratio of about 1.5 units to about 4 units of insulin glargine to 1 µg of AVE0010.

9. The pharmaceutical formulation of claim 8, wherein insulin glargine and AVE0010 are present in a ratio of about 3 units of insulin glargine to 1 µg of AVE0010.

10. The pharmaceutical formulation of claim 8, wherein insulin glargine and AVE0010 are present in a ratio of about 2 units of insulin glargine to 1 µg of AVE0010.

11. A pharmaceutical formulation comprising a combination of:

(a) insulin glargine or a pharmaceutically acceptable salt thereof, and

(b) desPro³⁶exendin-4(1-39)-Lys₆-NH₂ (AVE0010) or a pharmaceutically acceptable salt thereof; wherein the dosage of insulin glargine is between about 5 units and about 100 units, and the dosage of AVE0010 is between about 5 µg and about 2 mg.

12. The pharmaceutical formulation of claim 11, wherein the dosage of insulin glargine is between about 15 units and about 80 units, and the dosage of AVE0010 is between about 10 µg and about 30 µg.

13. The pharmaceutical formulation of claim 11, wherein the dosage of insulin glargine is between about 15 units and about 80 units, and the dosage of AVE0010 is between about 10 µg and about 20 µg.

US 9,526,764 B2

29

14. The pharmaceutical formulation of claim 1, 8, or 11, wherein the pharmaceutical formulation is a liquid formulation suitable for parenteral administration.

15. The pharmaceutical formulation of claim 1, 8, or 11, wherein the pharmaceutical formulation is suitable for administration once daily.

* * * * *

30